



PCT/AU2004/001689

Australian Government

Patent Office
Canberra

I, JANENE PEISKER, TEAM LEADER EXAMINATION SUPPORT AND SALES hereby certify that annexed is a true copy of the Provisional specification in connection with Application No. 2003906680 for a patent by CYTOPIA PTY LTD as filed on 03 December 2003.



WITNESS my hand this
Twenty-second day of December 2004

A handwritten signature in black ink, appearing to read 'J. Peisker'.

JANENE PEISKER
TEAM LEADER EXAMINATION
SUPPORT AND SALES

BEST AVAILABLE COPY

141590721

AUSTRALIA
Patents Act 1990 (Cth)

PROVISIONAL SPECIFICATION

Cytopia Pty Ltd

Invention Title

Tubulin Inhibitors

The invention is described in the following statement:

Blake Dawson Waldron Patent Services
Level 39, 101 Collins Street
Melbourne VIC 3000
Telephone: + 61 3 9679 3065
Fax: + 61 3 9679 3111

Ref: 13555655

141390721

2.

Tubulin inhibitors

FIELD OF THE INVENTION

The present invention relates to methods of suppressing the growth of cancers and other proliferative diseases by the administration of compounds that act by binding to tubulin and to novel heterocyclic compounds suitable for such administration.

BACKGROUND OF THE INVENTION

There are many human and veterinary diseases that stem from processes of uncontrolled or abnormal cellular proliferation. Most important among these diseases is cancer, the generic name given to a broad range of cellular malignancies characterized by unregulated growth and lack of differentiation. Psoriasis is another disease that is characterized by uncontrolled or abnormal cellular proliferation. Psoriasis is a common chronic skin disease characterized by the presence of dry scales and plaques. The disease results from hyperproliferation of the epidermis and incomplete differentiation of keratinocytes. Psoriasis often involves the scalp, elbows, knees, back, buttocks, nails, eyebrows, and genital regions, and may range in severity from mild to extremely debilitating, resulting in psoriatic arthritis, pustular psoriasis, and exfoliative psoriatic dermatitis. There is, at present, no general therapeutic cure that exists for psoriasis. Whilst milder cases are often treated with topical corticosteroids, more severe cases may be treated with antiproliferative agents, such as the antimetabolite methotrexate, the DNA synthesis inhibitor hydroxyurea, and the microtubule disrupter colchicine.

Other diseases associated with an abnormally high level of cellular proliferation include restenosis, where vascular smooth muscle cells are involved; inflammatory disease states, where endothelial cells, inflammatory cells and glomerular cells are involved; myocardial infarction, where heart muscle cells are involved; glomerular nephritis, where kidney cells are involved; transplant rejection, where endothelial cells are involved; and infectious diseases such as HIV infection and malaria, where certain immune cells and/or other infected cells are involved.

141590721

3.

Inhibition of cellular proliferation can be brought about by several mechanisms, including; alkylating agents; topoisomerase inhibitors; nucleotide analogues; antibiotics; hormone antagonists; and nucleic acid damaging agents; *inter alia*. One pharmacologically important mechanism of inhibiting cellular proliferation is by means of binding tubulin. Tubulin is an asymmetric dimer composed of alpha and beta subunits, that polymerizes to form structural components of the cytoskeleton called microtubules. Microtubules must be highly dynamic in order to carry out many of their functions. At certain stages of the cell cycle, or in particular cell types or organelles, stable microtubules are required, such as for transport within axons or for ciliary and flagellar movement.

Micro-tubules assemble during the G2 phase of the cell cycle, and participate in the formation of the mitotic spindle which facilitates the segregation of sister chromatids during the process of cell division. The essential role of microtubules in cell division and the ability of drugs that interact with tubulin to interfere with the cell cycle have made tubulin a successful target for applications that include anti-cancer drugs, fungicides, and herbicides. Typical tubulin ligands such as colchicine, paclitaxel, the *Vinca* alkaloids such as vinblastine, the epothilones, the halicondrins, benomyl and mebendazole directly inhibit cell division by binding to tubulin which leads to the arrest of the cell cycle at the G2/M boundary of mitosis. This mechanism is the basis of the therapeutic value of compounds of this type, such as treating gout with colchicine, restenosis with paclitaxel, cancer with paclitaxel, vinblastine, epothilones and halichondrins, and fungal infections with benomyl and malaria and helminths with mebendazole.

Interfering with microtubule dynamics or stability can inhibit cell division in several ways. Both stabilizing microtubules or inhibiting their polymerization will prevent the cytoskeleton restructuring that is required at several points in the cell cycle and lead to an arrest of the cell's progression from one stage in the cell cycle to the next. Three main classes of tubulin-binding drugs, namely colchicine analogues, *Vinca* alkaloids, and the taxanes, have been identified, each of which possesses a specific binding site on the β -tubulin molecule. Paclitaxel (TaxolTM) and related taxanes represent a class of drugs that stabilize microtubules, a process that ultimately leads to the "freezing" of the microtubule structures so that they cannot be restructured (Jordan M A. and Wilson L., 1998). Subsequent arrest at mitosis induces the apoptotic mechanism to cause cell death. A number of colchicine analogues, as well as several other compounds that bind to the same site on β -tubulin as colchicine disrupt tubulin polymerization and disrupt microtubular

141590721

4.

formation. Vinblastine and several other vinca-related drugs bind to a site that is distinct from the colchicine site. Compounds that bind at the Vinca-site prevent microtubule formation and destabilize microtubules (Jordan et al, 1986; Rai and Wolff (1996). This invention is therefore directed to compounds that potentially modulate microtubule dynamics by binding to tubulin.

Accordingly, the present invention aims to provide compounds which are directly or indirectly toxic to actively dividing cells and are useful in the treatment of cancer, viral and bacterial infections, vascular restenosis, inflammatory diseases, autoimmune diseases, or psoriasis. The present invention is also directed to therapeutic compositions for treating said conditions. Further aspects of the invention are to provide methods for killing actively proliferating cells, such as cancerous, bacterial, or epithelial cells, and treating all types of cancers, infections, inflammatory, and generally proliferative conditions. A further aspect relates to provide methods for treating other medical conditions characterized by the presence of rapidly proliferating cells, such as psoriasis and other skin disorders.

In one embodiment, the method of the invention is used in the treatment of sarcomas, carcinomas and/or leukemias. Exemplary disorders for which the subject method can be used alone or as part of a treatment regimen include: fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilms' tumor, cervical cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, meningioma, melanoma, neuroblastoma, and retinoblastoma.

In certain embodiments, the method of the invention is to be used to treat disorders such as carcinomas forming from tissue of the breast, prostate, kidney, bladder or colon.

141590721

5.


In other embodiments, the method of the invention is used to treat hyperplastic or neoplastic disorders arising in adipose tissue, such as adipose cell tumors, e.g., lipomas, fibrolipomas, lipoblastomas, lipomatosis, hibemomas, hemangiomas and/or liposarcomas.

- 5 In still other embodiments, infectious and parasitic agents (e.g. bacteria, trypanosomes, fungi, etc) can also be controlled using the subject compositions and compounds.

SUMMARY OF THE INVENTION

- 10 The present inventors have found that a group of compounds based upon a substituted heterocyclic scaffold inhibit the growth and proliferation of cancer cells. The present inventors have further shown that these compounds can bind to tubulin. Such compounds would also be useful in the treatment of other hyperproliferation related disorders.

- 15 Accordingly, in a first aspect the present invention provides a method of modulating microtubule polymerisation in a subject, the method comprising administering a therapeutically effective amount of at least one compound of the general formula



I

or pharmaceutically acceptable prodrugs, salts, hydrates, solvates, crystal forms or diastereomers thereof, wherein:

- 20 R^1 is H, C_{1-4} alkyl;

Q is a bond, or C_{1-4} alkyl;

A is aryl, hetaryl optionally substituted with 0-3 substituents independently chosen from halogen, C_{1-4} alkyl, CH_2F , CHF_2 , CF_3 , CN, aryl, hetaryl, OCF_3 ,

141590721

6.

OC₁₋₄alkyl, OC₂₋₅alkyl\NR4R5, Oaryl, Ohetaryl, CO₂R4, CONR4R5, nitro, NR4R5,
C₁₋₄alkyl\NR4R5, NR6C₁₋₄alkyl\NR4R5, NR4COR5, NR6CONR4R5, NR4SO₂R5;

5 R4, R5 are each independently H, C₁₋₄alkyl, C₁₋₄alkyl cycloalkyl, C₁₋₄alkyl
cyclohetalkyl, aryl, hetaryl, C₁₋₄alkyl aryl, C₁₋₄alkyl hetaryl, or may be joined
to form an optionally substituted 3-8 membered ring optionally containing
an atom selected from O, S, NR7;

R6 is selected from H, C₁₋₄alkyl;

R7 is selected from H, C₁₋₄alkyl, aryl, hetaryl, C₁₋₄alkyl aryl, C₁₋₄alkyl
hetaryl;

10 R2 is 0-2 substituents independently selected from halogen, C₁₋₄alkyl, OH,
OC₁₋₄alkyl, CH₂F, CHF₂, CF₃, OCF₃, CN, C₁₋₄alkyl\NR8R9, OC₁₋₄alkyl\NR8R9, CO₂R8,
CONR8R9, NR8R9, NR8COR9, NR10CONR8R9, NR8SO₂R9;

15 R8, R9 are each independently H, C₁₋₄alkyl, C₁₋₄alkyl cycloalkyl, C₁₋₄alkyl
cyclohetalkyl, aryl, hetaryl, C₁₋₄alkyl aryl, C₁₋₄alkyl hetaryl, or may be joined
to form an optionally substituted 3-8 membered ring optionally containing
an atom selected from O, S, NR11;

R10 is selected from H, C₁₋₄alkyl, aryl or hetaryl;

R11 is selected from H, C₁₋₄alkyl, aryl, hetaryl, C₁₋₄alkyl aryl, C₁₋₄alkyl
hetaryl;

20 Y is halogen, OH, NR12R13, NR14COR12, NR14CONR12R13, N14SO₂R13;

25 R12 and R13 are each independently H, CH₂F, CHF₂, CF₃, CN, C₁₋₄alkyl
optionally substituted with OH, OC₁₋₄alkyl or NR15R16, cycloalkyl;
cyclohetalkyl, C₁₋₄alkyl cycloalkyl, C₁₋₄alkyl cyclohetalkyl, or may be joined
to form an optionally substituted 3-6 membered ring optionally containing
an atom selected from O, S, NR14

R14, R15 and R16 are each independently selected from H, C₁₋₄alkyl;

n = 0-4;

7.

5

R17 is selected from H, C₁₋₄ alkyl.

II

Q is a bond, or C₁₋₄ alkyl;

15

20

R6 is selected from H, C₁₋₄ alkyl;

141590721

8.




R7 is selected from H, C₁₋₄ alkyl, aryl, hetaryl, C₁₋₄ alkyl aryl, C₁₋₄ alkyl hetaryl;

R2 is 0-2 substituents independently selected from C₁₋₄alkyl and OC₁₋₄alkyl;






5 Y is CH₂OH, OC₁₋₄alkylOH, OC₁₋₄alkylR12, OC₁₋₄alkylNR12NR13, C(O)R12, CH₂R12, COOR12, CONR12R13, OCONR12R13, CH₂NR12R13, NHCOR12, NHCONR12R13,

R12 and R13 are each independently H, C₁₋₂ alkyl, (CH₂)₃NEt₂, (CH₂)₂NMe₂, (CH₂)₂NH₂, (CH₂)₂OH,



10



15




20




141590721

9.



and



W is selected from H, C_{1-4} alkyl, C_{2-6} alkenyl; where C_{1-4} alkyl or C_{2-6} alkenyl may be optionally substituted with C_{1-4} alkyl, OH, OC_{1-4} alkyl, NR15R16;

R15, and R16 are each independently H, C_{1-4} alkyl, C_{1-4} alkyl cycloalkyl, C_{1-4} alkyl cycloheptalkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR17


R17 is selected from H, C_{1-4} alkyl;

wherein when Y is CH_2R_{12} then R12 is not H, C_{1-2} alkyl.

In a third aspect, the present invention provides a composition comprising a carrier and at least one compound of the second aspect.

In a fourth aspect the present invention provides a method of treating a hyperproliferation-related disorder in a subject, the method comprising administering a therapeutically effective amount of at least one compound of the second aspect or a composition of the third aspect.

In a fifth aspect the present invention provides a compound of the general formula



III

20

141590721

10.

or pharmaceutically acceptable prodrugs, salts, hydrates, solvates, crystal forms or diastereomers thereof, wherein:

X_1, X_2, X_3, X_4 are selected from the following:

- 5 (i) X_1 and X_2 are N and X_3 and X_4 are C independently substituted with Y;
(ii) X_1 and X_4 are N and X_2 and X_3 are C independently substituted with Y;
(iii) X_1 and X_3 are N and X_2 and X_4 are C independently substituted with Y;
(iv) X_2 and X_4 are N and X_1 and X_3 are C independently substituted with Y;
(v) X_1 is N and X_2, X_3 , and X_4 are C independently substituted with Y;
10 (vi) X_3 is N and X_1, X_2 , and X_4 are C independently substituted with Y;
(vii) X_4 is N and X_1, X_2 , and X_3 are C independently substituted with Y;
(viii) X_2 is N and X_1, X_3 , and X_4 are C independently substituted with Y; and
(ix) X_1, X_2 and X_3 are N and X_4 is C substituted with Y;

15 R1 is H, C_{1-6} alkyl, C_{1-6} alkylNR5R6, C_{1-6} alkylNR5COR6, C_{1-6} alkylNR5SO₂R6,
 C_{1-6} alkylCO₂R5, C_{1-6} alkylCONR5R6;

R5 and R6 are each independently H, C_{1-4} alkyl, aryl, hetaryl, C_{1-4} alkylaryl,
 C_{1-4} alkylhetaryl or may be joined to form an optionally substituted 3-8
membered ring optionally containing an atom selected from O, S, NR7;

R7 is selected from H, C_{1-4} alkyl;

20 R2 is selected from C_{1-6} alkylOH, OC_{2-6} alkylOH, C_{1-6} alkylNR8R9, OC_{2-6} alkylNR8R9,
 C_{1-6} alkylNR8COR9, OC_{2-6} alkylNR8COR9, C_{1-6} alkylhetaryl, OC_{2-6} alkylhetaryl,
OCONR8R9, NR8COOR9, NR10CONR8R9, CONR8R9, NR8COR12;

141590721

11.

R8, R9 are each independently H, C₁₋₄alkyl, C₁₋₄alkylNR₁₁R₁₃, hetaryl, cyclohetalkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR₁₄;

R₁₂ is C₂₋₄alkyl, C₁₋₄alkylNR₁₁R₁₃, hetaryl, cyclohetalkyl;

5

R₁₁, R₁₃ are each independently H, C₁₋₄alkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR₁₄;

R₁₄ is selected from H, C₁₋₄alkyl;

R₁₀ is H, C₁₋₄alkyl;

10

R₃ and R₄ are each independently H, halogen, C₁₋₄alkyl, OH, OC₁₋₄alkyl, CF₃, OCF₃;

Q is a bond, or C₁₋₄alkyl;

W is selected from H, C₁₋₄alkyl, C₂₋₄alkenyl; where C₁₋₄alkyl or C₂₋₄alkenyl may be optionally substituted with C₁₋₄alkyl, OH, OC₁₋₄alkyl, NR₁₅R₁₆;

15

R₁₅, and R₁₆ are each independently H, C₁₋₄alkyl, C₁₋₄alkyl cycloalkyl, C₁₋₄alkyl cyclohetalkyl, aryl, hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR₁₇;

R₁₇ is selected from H, C₁₋₄alkyl;

20

A is aryl, hetaryl optionally substituted with 0-3 substituents independently chosen from halogen, C₁₋₄alkyl, CF₃, aryl, hetaryl, OCF₃, OC₁₋₄alkyl, OC₂₋₅alkylNR₁₈R₁₉, Oaryl, Ohetaryl, CO₂R₁₈, CONR₁₈R₁₉, NR₁₈R₁₉, C₁₋₄alkylNR₁₈R₁₉, NR₂₀C₁₋₄alkylNR₁₈R₁₉, NR₁₈COR₁₉, NR₂₀CONR₁₈R₁₉, NR₁₈SO₂R₁₉;

25

R₁₈, R₁₉ are each independently H, C₁₋₄alkyl, C₁₋₄alkyl cyclohetalkyl, aryl, hetaryl, C₁₋₄alkyl aryl, C₁₋₄alkyl hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR₂₁;

141590721

12.


R21 is selected from H, C₁₋₄alkyl;R20 is selected from H, C₁₋₄alkyl;Y is selected from H, C₁₋₄alkyl, OH, NR₂₂R₂₃;R22, R23 are each independently H, C₁₋₄alkyl.

- 5 In a sixth aspect, the present invention provides a composition comprising a carrier and one or more compounds of the fifth aspect.

- In a seventh aspect, the present invention provides a method of treatment of a hyperproliferation-related disorder in a subject, the method comprising administering a therapeutically effective amount of at least one of the compounds of the fifth aspect or a composition according to the sixth aspect.
- 10

In an eighth aspect, the present invention consists in a method of treating a protein-kinase related disorder in a subject, the method comprising administering a therapeutically effective amount of at least one of the compounds of the fifth aspect or a composition according to the sixth aspect.

- 15 In a ninth aspect, the present invention provides a compound of the general formula



V

- or pharmaceutically acceptable prodrugs, salts, hydrates, solvates, crystal forms or diastereomers thereof, wherein:
- 20

141590721

13.

X_1, X_2, X_3, X_4 are selected from the following:

- (i) X_1 and X_2 are N and X_3 and X_4 are C independently substituted with Y;
- (ii) X_1 and X_4 are N and X_2 and X_3 are C independently substituted with Y;
- (iii) X_2 and X_4 are N and X_1 and X_3 are C independently substituted with Y;
- (iv) X_1 is N and X_2, X_3 , and X_4 are C independently substituted with Y;
- (v) X_3 is N and X_1, X_2 , and X_4 are C independently substituted with Y;
- (vi) X_4 is N and X_1, X_2 , and X_3 are C independently substituted with Y;
- (vii) X_2 is N and X_1, X_3 , and X_4 are C independently substituted with Y; and
- (viii) X_1, X_2 and X_3 are N and X_4 is C substituted with Y;

10 R_1 is H, C_{1-6} alkyl, C_{1-6} alkylNR5R6, C_{1-6} alkylNR5COR6, C_{1-6} alkylNR5SO₂R6, C_{1-6} alkylCO₂R5, C_{1-6} alkylCONR5R6;

R_5 and R_6 are each independently H, C_{1-4} alkyl, aryl, hetaryl, C_{1-4} alkylaryl, C_{1-4} alkylhetaryl or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR7;

15 R_7 is selected from H, C_{1-4} alkyl;

R_2 is selected from OH, C_{1-6} alkylOH, OC_{2-6} alkylOH, C_{1-6} alkylNR8R9, OC_{2-6} alkylNR8R9, C_{1-6} alkylNR8COR9, OC_{2-6} alkylNR8COR9, C_{1-6} alkylhetaryl, OC_{2-6} alkylhetaryl, $OCONR8R9$, $NR8COOR9$, $NR10CONR8R9$, $CONR8R9$, $NR8COR12$;

20 R_8, R_9 are each independently H, C_{1-4} alkyl, C_{1-4} alkylNR11R13, hetaryl, cyclohetalkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR14;

R_{12} is C_{2-4} alkyl, C_{1-4} alkylNR11R13, hetaryl, cyclohetalkyl;

141590721

14.

R11, R13 are each independently H, C₁₋₄alkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR14;

R14 is selected from H, C₁₋₄alkyl;

5 R10 is H, C₁₋₄ alkyl;

R3 and R4 are each independently H, halogen, C₁₋₄alkyl, OH, OC₁₋₄alkyl, CF₃, OCF₃;

Q is a bond, or C₁₋₄alkyl;

W is selected from H, C₁₋₄alkyl, C₂₋₆alkenyl; where C₁₋₄alkyl or C₂₋₆alkenyl may be optionally substituted with C₁₋₄alkyl, OH, OC₁₋₄alkyl, NR15R16;

10 R15, and R16 are each independently H, C₁₋₄alkyl, C₁₋₄alkyl cycloalkyl, C₁₋₄alkyl cyclohetalkyl, aryl, hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR17;

R17 is selected from H, C₁₋₄alkyl;

15 A is aryl, hetaryl optionally substituted with 0-3 substituents independently chosen from halogen, C₁₋₄ alkyl, CF₃, aryl, hetaryl, OCF₃, OC₁₋₄alkyl, OC₂₋₃alkylNR18R19, Oaryl, Ohetaryl, CO₂R18, CONR18R19, NR18R19, C₁₋₄alkylNR18R19, NR20C₁₋₄alkylNR18R19, NR18COR19, NR20CONR18R19, NR18SO₂R19;

20 R18, R19 are each independently H, C₁₋₄ alkyl, C₁₋₄ alkyl cyclohetalkyl, aryl, hetaryl, C₁₋₄alkyl aryl, C₁₋₄ alkyl hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR21;

R21 is selected from H, C₁₋₄ alkyl;

25 R20 is selected from H, C₁₋₄ alkyl;

Y is selected from H, C₁₋₄alkyl, OH, NR22R23;

141590721

15.

R22, R23 are each independently H, C₁₋₄ alkyl.


In a tenth aspect, the present invention provides a composition comprising a carrier and one or more compounds of the ninth aspect.

- 5 In an eleventh aspect, the present invention provides a method of treatment of a hyperproliferation-related disorder in a subject, the method comprising administering a therapeutically effective amount of at least one of the compounds of the ninth aspect or a composition according to the tenth aspect.

- 10 In a twelfth aspect, the present invention consists in a method of treating a protein-kinase related disorder in a subject, the method comprising administering a therapeutically effective amount of at least one of the compounds of the ninth aspect or a composition according to the tenth aspect.

DETAILED DESCRIPTION OF THE INVENTION

- 15 In a first aspect the present invention provides a method of modulating microtubule polymerisation in a subject, the method comprising administering a therapeutically effective amount of at least one compound of the general formula



I

or pharmaceutically acceptable salts, hydrates, solvates, crystal forms or diastereomers thereof, wherein:

- 20 R1 is H, C₁₋₄ alkyl;

Q is a bond, or C₁₋₄ alkyl;

A is aryl, hetaryl optionally substituted with 0-3 substituents independently chosen from halogen, C₁₋₄ alkyl, CH₂F, CHF₂, CF₃, CN, aryl, hetaryl, OCF₃,

141597721

16.

OC₁₋₄alkyl, OC₂₋₆alkylNR₄R₅, Oaryl, Ohetaryl, CO₂R₄, CONR₄R₅, nitro, NR₄R₅, C₁₋₄alkylNR₄R₅, NR₆C₁₋₄alkylNR₄R₅, NR₄COR₅, NR₆CONR₄R₅, NR₄SO₂R₅; and R₄, R₅ are each independently H, C₁₋₄alkyl, C₁₋₄alkyl cycloalkyl, C₁₋₄alkyl cyclohetalkyl, aryl, hetaryl, C₁₋₄alkyl aryl, C₁₋₄alkyl hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR₇; and R₆ is selected from H, C₁₋₄alkyl; and R₇ is selected from H, C₁₋₄alkyl, aryl, hetaryl, C₁₋₄alkyl aryl, C₁₋₄alkyl hetaryl;

R₂ is 0-2 substituents independently selected from halogen, C₁₋₄alkyl, OH, OC₁₋₄alkyl, CH₂F, CHF₂, CF₃, OCF₃, CN, C₁₋₄alkylNR₈R₉, OC₁₋₄alkylNR₈R₉, CO₂R₈, CONR₈R₉, NR₈R₉, NR₈COR₉, NR₁₀CONR₈R₉, NR₈SO₂R₉; and R₈, R₉ are each independently H, C₁₋₄alkyl, C₁₋₄alkyl cycloalkyl, C₁₋₄alkyl cyclohetalkyl, aryl, hetaryl, C₁₋₄alkyl aryl, C₁₋₄alkyl hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR₁₁; and R₁₀ is selected from H, C₁₋₄alkyl, aryl or hetaryl; and R₁₁ is selected from H, C₁₋₄alkyl, aryl, hetaryl, C₁₋₄alkyl aryl, C₁₋₄alkyl hetaryl;

Y is halogen; OH; NR₁₂R₁₃; NR₁₄COR₁₂; NR₁₄CONR₁₂R₁₃; N₁₄SO₂R₁₃; and R₁₂ and R₁₃ are each independently H; CH₂F; CHF₂; CF₃; CN; C₁₋₄alkyl optionally substituted with OH, OC₁₋₄alkyl or NR₁₅R₁₆; cycloalkyl; cyclohetalkyl; C₁₋₄alkyl cycloalkyl; C₁₋₄alkyl cyclohetalkyl; or may be joined to form an optionally substituted 3-6 membered ring optionally containing an atom selected from O, S, NR₁₄ and R₁₄, R₁₅ and R₁₆ are each independently selected from H, C₁₋₄alkyl;

n = 0-4;

W is selected from H, C₁₋₄alkyl, C₂₋₆alkenyl; where C₁₋₄alkyl or C₂₋₆alkenyl may be optionally substituted with C₁₋₄alkyl, OH, OC₁₋₄alkyl, NR₁₅R₁₆; and R₁₅, and R₁₆ are each independently H, C₁₋₄alkyl, C₁₋₄alkyl cycloalkyl, C₁₋₄alkyl cyclohetalkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR₁₇ and R₁₇ is selected from H, C₁₋₄alkyl.


Preferably, the method is used in the treatment of a hyperproliferation-related disorder.

Preferably, the compound is selected from the compounds set out in Table 2.

141590721

17.

In a second aspect the present invention provides a compound of the general formula



II

or pharmaceutically acceptable prodrugs, salts, hydrates, solvates, crystal forms or diastereomers thereof, wherein:


R1 is H, C₁₋₄ alkyl;

Q is a bond, or C₁₋₄ alkyl;

A is aryl, hetaryl optionally substituted with 0-3 substituents independently chosen from halogen, C₁₋₄ alkyl, CH₂F, CHF₂, CF₃, CN, aryl, hetaryl, OCF₃, OC₁₋₄alkyl, OC₂₋₆alkylNR₄R₅, Oaryl, Ohetaryl, CO₂R₄, CONR₄R₅, nitro, NR₄R₅, C₁₋₄ alkylNR₄R₅, NR₆C₁₋₄alkylNR₄R₅, NR₄COR₅, NR₆CONR₄R₅, NR₄SO₂R₅; and R₄, R₅ are each independently H, C₁₋₄ alkyl, C₁₋₄ alkyl cycloalkyl, C₁₋₄ alkyl cyclohetalkyl, aryl, hetaryl, C₁₋₄alkyl aryl, C₁₋₄ alkyl hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR₇; and R₆ is selected from H, C₁₋₄ alkyl; and R₇ is selected from H, C₁₋₄ alkyl, aryl, hetaryl, C₁₋₄ alkyl aryl, C₁₋₄ alkyl hetaryl;

R₂ is 0-2 substituents independently selected from C₁₋₄alkyl and OC₁₋₄alkyl;





Y is CH₂OH, OC₁₋₄alkylOH, OC₁₋₄alkylR₁₂, OC₁₋₄alkylNR₁₂NR₁₃, C(O)R₁₂, CH₂R₁₂, COOR₁₂, CONR₁₂R₁₃, OCONR₁₂R₁₃, CH₂NR₁₂R₁₃, NHCOR₁₂, NHCONR₁₂R₁₃, and R₁₂ and R₁₃ are each independently H, C₁₋₂ alkyl, (CH₂)₃NEt₂, (CH₂)₂NMe₂, (CH₂)₃NH₂, (CH₂)₂OH,





141590721

18.


5




10



15



and

 $n = 0-4$;

20

W is selected from H, C_{1-4} alkyl, C_{2-6} alkenyl; where C_{1-4} alkyl or C_{2-6} alkenyl may be optionally substituted with C_{1-4} alkyl, OH, OC_{1-4} alkyl, NR15R16; and R15, and R16 are each independently H, C_{1-4} alkyl, C_{1-4} alkyl cycloalkyl, C_{1-4} alkyl cyclohetalkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR17 and R17 is selected from H, C_{1-4} alkyl;

wherein when Y is CH_2 , R12 then R12 is not H, C_{1-4} alkyl.

141590721


19.

In a third aspect, the present invention comprises a composition comprising a carrier and at least one compound of the second aspect.

In a fourth aspect the present invention comprises a method of treating a hyperproliferation-related disorder in a subject, the method comprising administering a therapeutically effective amount of at least one compound of the second aspect or a composition of the third aspect.

Preferably, the hyperproliferation-related disorder is treatable by the modulation of microtubule polymerisation.

In a fifth aspect the present invention provides a compound of the general formula



III

or pharmaceutically acceptable prodrugs, salts, hydrates, solvates, crystal forms or diastereomers thereof, wherein:

X_1, X_2, X_3, X_4 are selected from the following:

- (i) X_1 and X_2 are N and X_3 and X_4 are C independently substituted with Y;
- (ii) X_1 and X_4 are N and X_2 and X_3 are C independently substituted with Y;
- (iii) X_1 and X_3 are N and X_2 and X_4 are C independently substituted with Y;
- (iv) X_2 and X_4 are N and X_1 and X_3 are C independently substituted with Y;

141590721

20.

- (v) X_1 is N and X_2 , X_3 , and X_4 are C independently substituted with Y;
(vi) X_3 is N and X_1 , X_2 , and X_4 are C independently substituted with Y;
(vii) X_4 is N and X_1 , X_2 , and X_3 are C independently substituted with Y;
(viii) X_2 is N and X_1 , X_3 , and X_4 are C independently substituted with Y; and
(ix) X_1 , X_2 and X_3 are N and X_4 is C substituted with Y;

R1 is H, C_{1-6} alkyl, C_{1-6} alkylNR5R6, C_{1-6} alkylNR5COR6, C_{1-6} alkylNR5SO₂R6, C_{1-6} alkylCO₂R5, C_{1-6} alkylCONR5R6, where R5 and R6 are each independently H, C_{1-4} alkyl, aryl, hetaryl, C_{1-4} alkylaryl, C_{1-4} alkylhetaryl or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR7;

R7 is selected from H, C_{1-4} alkyl;

R2 is selected from C_{1-4} alkylOH, OC_{2-6} alkylOH, C_{1-6} alkylNR8R9, OC_{2-6} alkylNR8R9, C_{1-6} alkylNR8COR9, OC_{2-6} alkylNR8COR9, C_{1-6} alkylhetaryl, OC_{2-6} alkylhetaryl, $OCONR8R9$, $NR8COOR9$, $NR10CONR8R9$, $CONR8R9$, $NR8COR12$;

R8, R9 are each independently H, C_{1-4} alkyl, C_{1-4} alkylNR11R13, hetaryl, cyclohetalkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR14;

R12 is C_{2-4} alkyl, C_{1-4} alkylNR11R13, hetaryl, cyclohetalkyl;

R11, R13 are each independently H, C_{1-4} alkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR14;

R14 is selected from H, C_{1-4} alkyl;

R10 is H, C_{1-4} alkyl;

R3 and R4 are each independently H, halogen, C_{1-4} alkyl, OH, OC_{1-4} alkyl, CF_3 , OCF_3 ;

Q is a bond, or C_{1-4} alkyl;

141590721

21.

W is selected from H, C₁₋₄alkyl, C₂₋₆alkenyl; where C₁₋₄alkyl or C₂₋₆alkenyl may be optionally substituted with C₁₋₄alkyl, OH, OC₁₋₄alkyl, NR15R16;

R15, and R16 are each independently H, C₁₋₄alkyl, C₁₋₄alkyl cycloalkyl, C₁₋₄alkyl cyclohetalkyl, aryl, hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR17;

R17 is selected from H, C₁₋₄alkyl;

A is aryl, hetaryl optionally substituted with 0-3 substituents independently chosen from halogen, C₁₋₄alkyl, CF₃, aryl, hetaryl, OCF₃, OC₁₋₄alkyl, OC₂₋₆alkylNR18R19, Oaryl, Ohetaryl, CO₂R18, CONR18R19, NR18R19, C₁₋₄alkylINR18R19, NR20C₁₋₄alkylINR18R19, NR18COR19, NR20CONR18R19, NR18SO₂R19;

R18, R19 are each independently H, C₁₋₄alkyl, C₁₋₄alkyl cyclohetalkyl, aryl, hetaryl, C₁₋₄alkyl aryl, C₁₋₄alkyl hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR21;

R21 is selected from H, C₁₋₄alkyl;

R20 is selected from H, C₁₋₄alkyl;

Y is selected from H, C₁₋₄alkyl, OH, NR22R23;

R22, R23 are each independently H, C₁₋₄alkyl.

In the above description it will be appreciated that:

C₁₋₄alkyl means an unsubstituted or optionally substituted straight or branched alkyl chain

Aryl means unsubstituted or optionally substituted phenyl or naphthyl.

Hetaryl means an unsubstituted or optionally substituted 5- or 6-membered heteroaromatic ring containing one or more heteroatoms selected from O, N, S.


141590721

22.

Cycloalkyl means a 3-8 membered saturated ring

Cyclohetalkyl means an optionally substituted 3-8 membered saturated ring containing 1-3 heteroatoms selected from O, S, NR₂₄, where R₂₄ is H, C₁₋₄ alkyl, aryl, hetaryl.

- 5 In a preferred embodiment the compound is selected from compounds of the general formula



IV

- 10 or pharmaceutically acceptable prodrugs, salts, hydrates, solvates, crystal forms or diastereomers thereof, wherein:

X₁, X₂, X₃, X₄ are selected from the following:

- 15 (i) X₁ and X₂ are N and X₃ and X₄ are C independently substituted with Y;
 (ii) X₁ and X₄ are N and X₂ and X₃ are C independently substituted with Y;
 (iii) X₁ and X₃ are N and X₂ and X₄ are C independently substituted with Y;
 (iv) X₂ and X₄ are N and X₁ and X₃ are C independently substituted with Y;
 (v) X₁ is N and X₂, X₃, and X₄ are C independently substituted with Y;
 (vi) X₃ is N and X₁, X₂, and X₄ are C independently substituted with Y;
 (vii) X₄ is N and X₁, X₂, and X₃ are C independently substituted with Y;
 (viii) X₂ is N and X₁, X₃, and X₄ are C independently substituted with Y; and

141590721

23.

(ix) X_1 , X_2 and X_3 are N and X_4 is C substituted with Y;

R1 is H, C_{1-4} alkyl, C_{1-4} alkylNR5R6, where R5 and R6 are each independently H, C_{1-4} alkyl, aryl, hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR7;

5 R7 is selected from H, C_{1-4} alkyl;

R2 is selected from C_{1-6} alkylOH, OC_{2-6} alkylOH, C_{1-6} alkylNR8R9, OC_{2-6} alkylNR8R9, C_{1-6} alkylNR8COR9, OC_{2-6} alkylNR8COR9, C_{1-6} alkylhetaryl, OC_{2-6} alkylhetaryl, OCONR8R9, NR8COOR9, NR10CONR8R9, CONR8R9, NR8COR12;

10 R8, R9 are each independently H, C_{1-4} alkyl, C_{1-4} alkylNR11R13, hetaryl, cyclohetalkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR14;

R12 is C_{2-4} alkyl, C_{1-4} alkylNR11R13, hetaryl, cyclohetalkyl;

15 R11, R13 are each independently H, C_{1-4} alkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR14;

R14 is selected from H, C_{1-4} alkyl;

R10 is H, C_{1-4} alkyl;

R3 and R4 are each independently H, halogen, C_{1-4} alkyl, OH, OC_{1-4} alkyl, CF_3 , OCF_3 ;

Q is CH;

20 W is selected from C_{1-4} alkyl, C_{2-6} alkenyl; where C_{1-4} alkyl or C_{2-6} alkenyl may be optionally substituted with C_{1-4} alkyl, OH, OC_{1-4} alkyl, NR15R16;

R15, and R16 are each independently H, C_{1-4} alkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR17;

25 R17 is selected from H, C_{1-4} alkyl;

141590721

24.

A is aryl, hetaryl optionally substituted with 0-2 substituents independently chosen from halogen, C₁₋₄ alkyl, CF₃, aryl, hetaryl, OCF₃, OC₁₋₄alkyl, OC₂₋₈alkylNR18R19, Oaryl, Ohetaryl, CO₂R18, CONR18R19, NR18R19, C₁₋₄alkylNR18R19, NR20C₁₋₄alkylNR18R19, NR18COR19, NR20CONR18R19, NR18SO₂R19;

R18, R19 are each independently H, C₁₋₄alkyl, C₁₋₄alkyl cyclohetalkyl, aryl, hetaryl, C₁₋₄alkyl aryl, C₁₋₄alkyl hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR21;

R21 is selected from H, C₁₋₄alkyl;

R20 is selected from H, C₁₋₄alkyl;

Y is selected from H, C₁₋₄alkyl, NR22R23;

R22, R23 are each independently H, C₁₋₄alkyl.

In the above description it will be appreciated that:

C₁₋₄ alkyl means an unsubstituted or optionally substituted straight or branched alkyl chain

Aryl means unsubstituted or optionally substituted phenyl or naphthyl.

Hetaryl means an unsubstituted or optionally substituted 5- or 6-membered heteroaromatic ring containing one or more heteroatoms selected from O, N, S.

Cycloalkyl means a 3-8 membered saturated ring

Cyclohetalkyl means an optionally substituted 3-8 membered saturated ring containing 1-3 heteroatoms selected from O, S, NR24, where R24 is H, C₁₋₄ alkyl, aryl, hetaryl.

In a sixth aspect, the present invention provides a composition comprising a carrier and one or more compounds of the fifth aspect.

141590721


25.

In a seventh aspect, the present invention provides a method of treatment of a hyperproliferation-related disorder in a subject, the method comprising administering a therapeutically effective amount of at least one of the compounds of the fifth aspect or a composition according to the sixth aspect.

- 5 Preferably, the hyperproliferation-related disorder is treatable by the modulation of microtubule polymerisation.

- In an eighth aspect, the present invention consists in a method of treating a protein-kinase related disorder in a subject, the method comprising administering a therapeutically effective amount of at least one of the compounds of the fifth aspect or a composition
10 according to the sixth aspect.

In a ninth aspect, the present invention provides a compound of the general formula



V

- 15 or pharmaceutically acceptable prodrugs, salts, hydrates, solvates, crystal forms or diastereoisomers thereof, wherein:

X_1, X_2, X_3, X_4 are selected from the following:

- (i) X_1 and X_2 are N and X_3 and X_4 are C independently substituted with Y;
(ii) X_1 and X_4 are N and X_2 and X_3 are C independently substituted with Y;
20 (iii) X_2 and X_4 are N and X_1 and X_3 are C independently substituted with Y;

141590721

26.

- (iv) X_1 is N and X_2 , X_3 , and X_4 are C independently substituted with Y;
(v) X_3 is N and X_1 , X_2 , and X_4 are C independently substituted with Y;
(vi) X_4 is N and X_1 , X_2 , and X_3 are C independently substituted with Y;
(vii) X_2 is N and X_1 , X_3 , and X_4 are C independently substituted with Y; and
5 (viii) X_1 , X_2 and X_3 are N and X_4 is C substituted with Y;

R1 is H, C_{1-6} alkyl, C_{1-6} alkylNR5R6, C_{1-6} alkylNR5COR6, C_{1-6} alkylNR5SO₂R6, C_{1-6} alkylCO₂R5, C_{1-6} alkylCONR5R6, where R5 and R6 are each independently H, C_{1-6} alkyl, aryl, hetaryl, C_{1-4} alkylaryl, C_{1-4} alkylhetaryl or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected
10 from O, S, NR7;

R7 is selected from H, C_{1-4} alkyl;

R2 is selected from OH, C_{1-6} alkylOH, OC₂₋₆alkylOH, C_{1-6} alkylNR8R9, OC₂₋₆alkylNR8R9, C_{1-6} alkylNR8COR9, OC₂₋₆alkylNR8COR9, C_{1-6} alkylhetaryl, OC₂₋₆alkylhetaryl, OCONR8R9, NR8COOR9, NR10CONR8R9, CONR8R9,
15 NR8COR12;

R8, R9 are each independently H, C_{1-4} alkyl, C_{1-4} alkylNR11R13, hetaryl, cyclohetalkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR14;

R12 is C_{2-4} alkyl, C_{1-6} alkylNR11R13, hetaryl, cyclohetalkyl;

20 R11, R13 are each independently H, C_{1-4} alkyl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR14;

R14 is selected from H, C_{1-4} alkyl;

R10 is H, C_{1-4} alkyl;

25 R3 and R4 are each independently H, halogen, C_{1-4} alkyl, OH, OC₁₋₄alkyl, CF₃, OCF₃;

141590721

27.

Q is a bond, or C₁₋₄alkyl;

W is selected from H, C₁₋₄alkyl, C₂₋₆alkenyl; where C₁₋₄alkyl or C₂₋₆alkenyl may be optionally substituted with C₁₋₄alkyl, OH, OC₁₋₄alkyl, NR15R16;

5 R15, and R16 are each independently H, C₁₋₄alkyl, C₁₋₄alkyl cycloalkyl, C₁₋₄alkyl cyclohetalkyl, aryl, hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR17;

R17 is selected from H, C₁₋₄alkyl;

10 A is aryl, hetaryl optionally substituted with 0-3 substituents independently chosen from halogen, C₁₋₄alkyl, CF₃, aryl, hetaryl, OCF₃, OC₁₋₄alkyl, OC₂₋₆alkyl, NR18R19, Oaryl, Ohetaryl, CO₂R18, CONR18R19, NR18R19, C₁₋₄alkyl, NR20C₁₋₄alkyl, NR18COR19, NR20CONR18R19, NR18SO₂R19;

15 R18, R19 are each independently H, C₁₋₄alkyl, C₁₋₄alkyl cyclohetalkyl, aryl, hetaryl, C₁₋₄alkyl aryl, C₁₋₄alkyl hetaryl, or may be joined to form an optionally substituted 3-8 membered ring optionally containing an atom selected from O, S, NR21;

R21 is selected from H, C₁₋₄alkyl;

R20 is selected from H, C₁₋₄alkyl;

20 Y is selected from H, C₁₋₄alkyl, OH, NR22R23;

R22, R23 are each independently H, C₁₋₄alkyl.

In the above description it will be appreciated that:

C₁₋₄alkyl means an unsubstituted or optionally substituted straight or branched alkyl chain

25 Aryl means unsubstituted or optionally substituted phenyl or naphthyl.

141590721

28.

Hetaryl means an unsubstituted or optionally substituted 5- or 6-membered heteroaromatic ring containing one or more heteroatoms selected from O, N, S.

Cycloalkyl means a 3-8 membered saturated ring

5 Cyclohetaryl means an optionally substituted 3-8 membered saturated ring containing 1-3 heteroatoms selected from O, S, NR²⁴, where R²⁴ is H, C₁₋₄ alkyl, aryl, hetaryl.

In a tenth aspect, the present invention provides a composition comprising a carrier and one or more compounds of the ninth aspect.

10 In an eleventh aspect, the present invention provides a method of treatment of a hyperproliferation-related disorder in a subject, the method comprising administering a therapeutically effective amount of at least one of the compounds of the ninth aspect or a composition according to the tenth aspect.

Preferably, the hyperproliferation-related disorder is treatable by the modulation of microtubule polymerisation.

15 In a twelfth aspect, the present invention consists in a method of treating a protein-kinase related disorder in a subject, the method comprising administering a therapeutically effective amount of at least one of the compounds of the ninth aspect or a composition according to the tenth aspect.

20 The compounds of this invention include all conformational isomers (eg. cis and trans isomers). The compounds of the present invention have asymmetric centers and therefore exist in different enantiomeric and diastereomeric forms. This invention relates to the use of all optical isomers and stereoisomers of the compounds of the present invention, and mixtures thereof, and to all pharmaceutical compositions and methods of treatment that may employ or contain them. The compounds of formula I, II, III, IV and V may also exist
25 as tautomers. This invention relates to the use of all such tautomers and mixtures thereof.

Compounds of formula I, II, III, IV and V having free amino, amido, hydroxy or carboxylic groups can be converted into prodrugs. Prodrugs include compounds wherein an amino acid residue, or a polypeptide chain of two or more (eg. two, three or four) amino acid residues which are covalently joined through peptide bonds to free amino,

141590721

29.

hydroxy and carboxylic acid groups of compounds of formulae I, II, III, IV and V. The amino acid residues include the 20 naturally occurring amino acids commonly designated by three letter symbols and also include, 4-hydroxyproline, hydroxylysine, demosine, isodemossine, 3-methylhistidine, norvaline, beta-alanine, gamma-aminobutyric acid, citrulline, homocysteine, homoserine, ornithine and methionine sulfone. Prodrugs also include compounds wherein carbonates, carbamates, amides and alkyl esters which are covalently bonded to the above substituents of formulae I, II, III, IV and V through the carbonyl carbon prodrug sidechain. Prodrugs also include phosphate derivatives of compounds of formulae I, II, III, IV and V (such as acids, salts of acids, or esters) joined through a phosphorus-oxygen bond to a free hydroxyl of compounds of formulae I, II, III, IV and V. Prodrugs also include compounds wherein acyloxyalkyl or phosphonooxyalkyl moieties are covalently attached to compounds of formula I, II, III, IV and V possessing a free hydroxyl group. Acyloxyalkyl or phosphonooxyalkyl moieties may also be covalently attached to compounds of formulae I, II, III, IV and V possessing a pyridyl ring through formation of a N-(acyloxyalkyl)- or N-(phosphonooxyalkyl)-pyridinium salt. This invention also encompasses pharmaceutical compositions containing prodrugs of compounds of the formula I, II, III, IV and V. In addition, this invention encompasses methods of treating or preventing disorders that can be treated or prevented by the modulation of microtubule dynamics comprising administering prodrugs of compounds of the formula I, II, III, IV and V. This invention further encompasses methods of treating or preventing disorders that can be treated or prevented by the disruption of microtubules comprising administering prodrugs of compounds of the formula II, III, IV and V.

In a still further preferred embodiment the compound possesses *S* chirality at the chiral carbon bearing W, where W is C₁₋₄ alkyl or C₁₋₄ alkylamino. The compound can be used as a purified isomer or as a mixture of any ratio of isomers. It is however preferred that the mixture comprises at least 70%, 80%, 90%, 95%, or 99% of the preferred isomer.

In a yet further preferred embodiment, the present invention provides a compound selected from the compounds of Table 1; compositions comprising a carrier and one or more compounds of Table 1; and, a method of treatment of a hyperproliferation-related disorder in a subject, the method comprising administering at least one of the compounds of Table 1.

141590721

30.

More preferably, the hyperproliferation related disorder is treatable by the modulation of microtubule polymerisation.

The present invention further provides for a method of modulating microtubule polymerisation in a subject, the method comprising the administration of a
5 therapeutically effective amount of a compound of any one of Formulas I, II, III, IV and V.

In a preferred embodiment, there is provided a method of modulating microtubule polymerisation in a subject, the method comprising the administration of a therapeutically effective amount of a compound selected from the compounds of Table 1.

The present invention provides that the compounds of formula I, II, III, IV and V may be
10 administered for the treatment of hyperproliferation-related disorders. Preferably, the hyperproliferation-related disorder is treatable by the modulation of microtubule polymerisation. In a preferred embodiment, the hyperproliferation-related disorder or disease state is selected from the group consisting of Cancer, such as fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma,
15 angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilms' tumor, cervical cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma,
25 meningioma, melanoma, neuroblastoma, and retinoblastoma, and carcinomas forming from tissue of the breast, prostate, kidney, bladder or colon, and neoplastic disorders arising in adipose tissue, such as adipose cell tumors, e.g., lipomas, fibrolipomas, lipoblastomas, lipomatosis, hibemomas, hemangiomas and/or liposarcomas; infectious diseases such as viral, malarial and bacterial infections; vascular restenosis; inflammatory
30 diseases, such as autoimmune diseases, glomerular nephritis myocardial infarction and psoriasis.

141590721

31.

The compounds of formula II, II and IV may also be used in the treatment of a hyperproliferation-related disorder or disease state and/or a protein-kinase related disorder or disease state. Preferably, the disease state is selected from the group consisting of Atopy, such as Allergic Asthma, Atopic Dermatitis (Eczema), and Allergic Rhinitis; Cell Mediated Hypersensitivity, such as Allergic Contact Dermatitis and Hypersensitivity Pneumonitis; Rheumatic Diseases, such as Systemic Lupus Erythematosus (SLE), Rheumatoid Arthritis, Juvenile Arthritis, Sjögren's Syndrome, Scleroderma, Polymyositis, Ankylosing Spondylitis, Psoriatic Arthritis; Other autoimmune diseases such as Type I diabetes, autoimmune thyroid disorders, and Alzheimer's disease; Viral Diseases, such as Epstein Barr Virus (EBV), Hepatitis B, Hepatitis C, HIV, HTLV 1, Varicella-Zoster Virus (VZV), Human Papilloma Virus (HPV).

In the treatment or prevention of hyperproliferation-related disorders and/or protein kinase related disorders an appropriate dosage level will generally be about 0.01 to 500 mg per kg patient body weight per day which can be administered in single or multiple doses. Preferably, the dosage level will be about 0.1 to about 250 mg/kg per day; more preferably about 0.5 to about 100 mg/kg per day. A suitable dosage level may be about 0.01 to 250 mg/kg per day, about 0.05 to 100 mg/kg per day, or about 0.1 to 50 mg/kg per day. Within this range the dosage may be 0.05 to 0.5, 0.5 to 5 or 5 to 50 mg/kg per day. For oral administration, the compositions are preferably provided in the form of tablets containing 1.0 to 1000 milligrams of the active ingredient, particularly 1.0, 5.0, 10.0, 15.0, 20.0, 25.0, 50.0, 75.0, 100.0, 150.0, 200.0, 250.0, 300.0, 400.0, 500.0, 600.0, 750.0, 800.0, 900.0, and 1000.0 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated. The compounds may be administered on a regimen of 1 to 4 times per day, preferably once or twice per day.

It will be understood, however, that the specific dose level and frequency of dosage for any particular patient may be varied and will depend upon a variety of factors including the activity of the specific compound employed, the metabolic stability and length of action of that compound, the age, body weight, general health, sex, diet, mode and time of administration, rate of excretion, drug combination, the severity of the particular condition, and the host undergoing therapy.

In addition to primates, such as humans, a variety of other mammals can be treated according to the method of the present invention. For instance, mammals including, but

141590721

32.

not limited to, cows, sheep, goats, horses, dogs, cats, guinea pigs, rats or other bovine, ovine, equine, canine, feline, rodent or murine species can be treated. However, the method can also be practiced in other species, such as avian species (e.g., chickens).

5 Diseases and conditions associated with inflammation and infection can be treated using the method of the present invention. In a preferred embodiment, the disease or condition is one in which the actions of eosinophils and/or lymphocytes are to be inhibited or promoted, in order to modulate the inflammatory response.

10 The subjects treated in the above methods, in whom which cell growth inhibition is desired, are mammals, including, but not limited to, cows, sheep, goats, horses, dogs, cats, guinea pigs, rats or other bovine, ovine, equine, canine, feline, rodent or murine species, and preferably a human being, male or female.

The term "therapeutically effective amount" means the amount of the subject composition that will elicit the biological or medical response of a tissue, system, animal or human that is being sought by the researcher, veterinarian, medical doctor or other clinician.

15 The terms "administration of" and or "administering a" compound should be understood to mean providing a compound of the invention to the individual in need of treatment.

20 The present invention provides pharmaceutical compositions comprising at least one of the compounds of the formula I, II, III, IV and V capable of treating a hyperproliferation-related disorder in an amount effective therefore, and a pharmaceutically acceptable vehicle or diluent. The compositions of the present invention may contain other therapeutic agents as described below, and may be formulated, for example, by employing conventional solid or liquid vehicles or diluents, as well as pharmaceutical additives of a type appropriate to the mode of desired administration (for example, excipients, binders, preservatives, stabilizers, flavors, etc.) according to techniques such as
25 those well known in the art of pharmaceutical formulation.

30 The compounds of the formula I, II, III, IV and V may be administered by any suitable means, for example, orally, such as in the form of tablets, capsules, granules or powders; sublingually; buccally; parenterally, such as by subcutaneous, intravenous, intramuscular, or intracisternal injection or infusion techniques (e.g., as sterile injectable aqueous or non-aqueous solutions or suspensions); nasally such as by inhalation spray; topically, such as

141590721

33.

in the form of a cream or ointment; or rectally such as in the form of suppositories; in dosage unit formulations containing non-toxic, pharmaceutically acceptable vehicles or diluents. The compounds may, for example, be administered in a form suitable for immediate release or extended release. Immediate release or extended release may be achieved by the use of suitable pharmaceutical compositions comprising the present compounds, or, particularly in the case of extended release, by the use of devices such as subcutaneous implants or osmotic pumps.

The term "composition" as used herein is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts. By "pharmaceutically acceptable" it is meant the carrier, diluent or excipient must be compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

The pharmaceutical compositions for the administration of the compounds of this invention may conveniently be presented in dosage unit form and may be prepared by any of the methods well known in the art of pharmacy. All methods include the step of bringing the active ingredient into association with the carrier which constitutes one or more accessory ingredients. In general, the pharmaceutical compositions are prepared by uniformly and intimately bringing the active ingredient into association with a liquid carrier or a finely divided solid carrier or both, and then, if necessary, shaping the product into the desired formulation. In the pharmaceutical composition the active object compound is included in an amount sufficient to produce the desired effect upon the process or condition of diseases. As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts.

The pharmaceutical compositions containing the active ingredient may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, or syrups or elixirs. Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of

141590721

34.

sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These excipients may be for example, inert

5 diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the

10 gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed. They may also be coated to form osmotic therapeutic tablets for control release.

Formulations for oral use may also be presented as hard gelatin capsules wherein the

15 active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin, or olive oil.

Aqueous suspensions contain the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for

20 example sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinyl-pyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long

25 chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more

30 preservatives, for example ethyl, or n-propyl, p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose or saccharin.

141590721

35.

Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavoring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present.

The pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally- occurring gums, for example gum acacia or gum tragacanth, naturally- occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example sorbitan monooleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening and flavoring agents.

Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative and flavoring and coloring agents.

The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleagenous suspension. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butane diol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In

141590721

36.

addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

5 The compounds of the present invention may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials are cocoa butter and polyethylene glycols.

10 For topical use, creams, ointments, jellies, solutions or suspensions, etc., containing the compounds of the present invention are employed. (For purposes of this application, topical application shall include mouthwashes and gargles.)

The compounds of the present invention can also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or
15 other lipid substances. Liposomes are formed by mono- or multilamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any non-toxic, physiologically acceptable and metabolisable lipid capable of forming liposomes can be used. The present compositions in liposome form can contain, in addition to a compound of the present invention, stabilisers, preservatives, excipients and the like. The preferred lipids
20 are the phospholipids and phosphatidyl cholines, both natural and synthetic. Methods to form liposomes are known in the art.

The pharmaceutical composition and method of the present invention may further comprise other therapeutically active compounds as noted herein which are usually applied in the treatment of the above mentioned pathological conditions. Selection of the
25 appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for
30 adverse side effects.

141590721

37.

Examples of other therapeutic agents include the following:

- Tyrosine kinase inhibitors, such as imatinib (Glivec™), and gefitinib (Iressa™) *inter alia*; cyclosporins (e.g., cyclosporin A), CTLA4-Ig, antibodies such as ICAM-3, anti-IL-2 receptor (Anti-Tac), anti-CD45RB, anti-CD2, anti-CD3 (OKT-3), anti-CD4, anti-CD80, anti-CD86,
- 5 agents blocking the interaction between CD40 and gp39, such as antibodies specific for CD40 and/or gp39 (i.e., CD154), fusion proteins constructed from CD40 and gp39 (CD40lg and CD8gp39), inhibitors, such as nuclear translocation inhibitors, of NF-kappa B function, such as deoxyspergualin (DSG), cholesterol biosynthesis inhibitors such as HMG CoA reductase inhibitors (lovastatin and simvastatin), non-steroidal
- 10 antiinflammatory drugs (NSAIDs) such as ibuprofen, aspirin, acetaminophen and cyclooxygenase inhibitors such as rofecoxib, steroids such as prednisolone or dexamethasone, gold compounds, antiproliferative agents such as methotrexate, FK506 (tacrolimus, Prograf), mycophenolate mofetil, antineoplastic agents such as azathioprine, VP-16, etoposide, fludarabine, cisplatin, doxorubicin, adriamycin, amsacrine,
- 15 camptothecin, cytarabine, gemcitabine, fluorodeoxyuridine, melphalan and cyclophosphamide, TNF- α inhibitors such as tenidap, anti-TNF antibodies or soluble TNF receptor, and rapamycin (sirolimus or Rapamune) or derivatives thereof.

- When other therapeutic agents are employed in combination with the compounds of the present invention they may be used for example in amounts as noted in the Physician
- 20 Desk Reference (PDR) or as otherwise determined by one of ordinary skill in the art.

- The pharmaceutical composition and method of the present invention may further comprise other therapeutically active compounds as noted herein which are known inhibitors or substrates of drug efflux systems or drug detoxification and excretory systems. Such systems include P-glycoprotein, multidrug resistance-associated protein,
- 25 lung resistance protein and glutathione S-transferase isoenzymes alpha, mu, pi, sigma, theta, zeta and kappa. Co-administration of drugs known to inhibit or reduce the activity of these systems may increase the efficacy of the compounds described in the present invention through increasing the amount of therapeutic agent in the cell. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages, thus
- 30 reducing the potential for adverse side effects. Examples of inhibitors or substrates for these systems include; verapamil, probenecid, dipyridamole, ethacrynic acid, indomethacin, sulfasalazine, buthionine sulfoximine, cyclosporin A and tamoxifen.

141590721

38.

Throughout this specification the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element, integer or step, or group of elements, integers or steps, but not the exclusion of any other element, integer or step, or group of elements, integers or steps.

- 5 All publications mentioned in this specification are herein incorporated by reference.

- Any discussion of documents, acts, materials, devices, articles or the like which has been included in the present specification is solely for the purpose of providing a context for the present invention. It is not to be taken as an admission that any or all of these matters form part of the prior art base or were common general knowledge in the field relevant to the present invention as it existed in Australia before the priority date of each claim of this application.
- 10

In order that the nature of the present invention may be more clearly understood preferred forms thereof will now be described by reference to the following non-limiting Examples.

141590721

39.

MATERIALS AND METHODS:

Compound Synthesis

Compounds are generally prepared in a 2-step process starting from a dihaloheterocycle.

5 The first step is a nucleophilic aromatic substitution to generate a monoamino-monohalo intermediate.

10 The nucleophilic aromatic substitution is typically carried out by addition of a primary or secondary amine to the di-halogenated heterocycle in a solvent such as ethanol, isopropanol, tert-butanol, dioxane, THF, DMF, ethoxyethanol, toluene or xylene. The reaction is typically performed at elevated temperature in the presence of excess amine or a non-nucleophilic base such as triethylamine or diisopropylethylamine, or an inorganic base such as potassium carbonate or sodium carbonate.

15 Alternatively, the amino substituent may be introduced through a transition metal catalysed amination reaction. Typical catalysts for such transformations include $\text{Pd}(\text{OAc})_2/\text{P}(\text{t-Bu})_3$, $\text{Pd}_2(\text{dba})_3/\text{BINAP}$ and $\text{Pd}(\text{OAc})_2/\text{BINAP}$. These reactions are typically carried out in solvents such as toluene or dioxane, in the presence of bases such as caesium carbonate or sodium or potassium tert-butoxide at temperatures ranging from room temperature to reflux.

20 The amines employed in the first step of the synthesis of these compounds are obtained commercially or are prepared using methods well known to those skilled in the art. Of particular interest are α -alkylbenzylamines which may be prepared through reduction of oximes. Typical reductants include lithium aluminium hydride, hydrogen gas in the presence of palladium on charcoal catalyst, Zn in the presence of hydrochloric acid, sodium borohydride in the presence of a Lewis acid such as TiCl_3 , ZrCl_4 , NiCl_2 and MoO_3 , or sodium borohydride in conjunction with Amberlyst H15 ion exchange resin and LiCl.

25 α -Alkylbenzylamines may also be prepared by reductive amination of the corresponding ketones. A classical method for such a transformation is the Leuckart-Wallach reaction, though catalytic conditions $(\text{HCO}_2\text{NH}_4, [(\text{CH}_3)_3\text{C}_2\text{RhCl}_2]_2)$ or alternative procedures (e.g. NH_4OAc , $\text{Na}(\text{CN})\text{BH}_2$) can also be used.

141590721

40.

α -Alkylbenzylamines may also be prepared from the corresponding α -alkylbenzyl alcohols. Such methods include derivatisation of the hydroxyl as a mesylate or tosylate and displacement with a nitrogen nucleophile, such as phthalimide or azide which is converted to the primary amine using conventional synthetic methods; or, displacement of the hydroxyl with a suitable nitrogen nucleophile under Mitsunobu-like conditions. α -Alkylbenzyl alcohols can be prepared by reduction of the corresponding ketones with a reducing agent such as sodium borohydride in a solvent such as methanol. Alternatively, α -alkylbenzyl alcohols can be obtained through addition of an alkyl metal species (such as a Grignard reagent) to a benzaldehyde derivative, typically performed at room temperature or below in solvents such as tetrahydrofuran.

α -Alkyl benzylamines of high optical purity may be prepared from chiral α -alkyl benzyl alcohols using the methods outlined above. The chiral α -alkyl benzyl alcohols may be obtained through chiral reduction of the corresponding ketones. Chiral reducing methods are now well known in organic chemistry and include enzymatic processes, asymmetric hydrogenation procedures and chiral oxazaborolidines.

The second step of the synthesis typically involves a palladium mediated cross-coupling of the monoamino-monochloro intermediate with a suitably functionalised coupling partner. Typical coupling partners are boronic acids or esters (Suzuki coupling: see for example Miyaura, N. and Suzuki, *Chem Rev.* 1995, 952457), stannanes (Stille coupling: see for example Stille, J.K., *Angew. Chem., Int. Ed. Engl.*, 1986, 25, 508), Grignard reagents (Kumada coupling: Kumada, M.; Tamao, K.; Sumitani, K. *Org. Synth.* 1988, Coll. Vol.6, 407.) or organozinc species (Negishi coupling: Negishi, E.; *J. Organomet. Chem.* 2002, 653, 34).

The Suzuki coupling is the preferred coupling method and is typically performed in a solvent such as DME, THF, DMF, ethanol, propanol, toluene, or 1,4-dioxane in the presence of a base such as potassium carbonate, lithium hydroxide, caesium carbonate, sodium hydroxide, potassium fluoride or potassium phosphate. The reaction may be carried out at elevated temperatures and the palladium catalyst employed may be selected from $\text{Pd}(\text{PPh}_3)_4$, $\text{Pd}(\text{OAc})_2$, $[\text{PdCl}_2(\text{dppf})]$, $\text{Pd}_2(\text{dba})_3/\text{P}(\text{t-Bu})_3$.

The products formed from this reaction sequence may be further derivatised using techniques well-known to those skilled in the art. Alternatively, derivatisation of the

141590721

41.


mono-amino mono-halo intermediate may be undertaken prior to displacement of the halo substituent. This derivatisation typically involves functionality originally present on the amine species and employs methods well known to those skilled in the art.

- 5 Alternatively the preparation sequence can be reversed beginning with the palladium mediated cross-coupling reaction to furnish a mono-halo heterocyclic species. Amine displacement of the halo substituent can then be undertaken using procedures outlined above.

Representative syntheses are reported below.


141590721

42.

Example 1**1-[4-(Trifluoromethyl)phenyl]butan-1-ol**

5 A 2M solution of propylmagnesium chloride in ether (4 ml, 8 mmol) was added to a solution of the aldehyde (1.14 g, 6.6 mmol) in dry THF (10 ml) cooled to 0 °C under N₂. The mixture was stirred for 16h at room temperature, after which time saturated ammonium chloride solution was added. The product was extracted into ethyl acetate, and the ethyl acetate layer dried and concentrated to furnish pure product (1.4g, 98%).

10 ¹H-n.m.r. (CDCl₃) δ 0.94 (t, 3H, J = 7.2Hz, CH₃), 1.41 (m, 2H, CH₂), 1.75 (m, CH₂, 2H), 4.77 (br s, 1H, CH), 7.44-7.62(m, 4H, ArH)

Example 2**1-(1-Azidobutyl)-4-(trifluoromethyl)benzene**


15 A solution of 1-[4-(trifluoromethyl)phenyl]butan-1-ol (1.4 g, 6.4 mmol) and diphenylphosphoryl azide (2.8 ml, 12.8 mmol) in THF (6 mL) cooled to -10 °C under N₂ was treated with DBU (1.9 mL, 12.8 mmol). The resulting solution was stirred at room temperature for 20 hours and then diluted with a mixture of ether and H₂O. The organic phase was dried and concentrated and the residue purified by column chromatography using hexane:ethyl acetate (10:1) as eluent to furnish pure azide (0.85 g, 54%).

20 ¹H-n.m.r. (CDCl₃) δ 0.94 (t, 3H, J = 7.2Hz, CH₃), 1.37 (m, 2H, CH₂), 1.75 (m, 2H, CH₂), 4.50 (t, 1H, CH), 7.42 (d, J = 7.8 Hz, 2H, ArH), 7.64 (d, 2H, J = 7.8Hz, ArH)

Example 3

141590721

43.

1-[4-(Trifluoromethyl)phenyl]butan-1-amine

A mixture of 1-(1-azidobutyl)-4-(trifluoromethyl)benzene (0.84 g, 3.5 mmol) and triphenylphosphine (1.8g, 6.9 mmol) in ethyl acetate (6 mL) and 10% HCl (6ml) was stirred at room temperature for 64 h. The aqueous phase was collected and the organic phase extracted with 10% HCl (3 x 5mL). The aqueous layers were combined and basified with 5M NaOH, and then extracted with ethyl acetate (5 x 15mL). The organic phase was dried and concentrated to give pure amine (0.4 g, 54%).

¹H-n.m.r. (CDCl₃) δ 0.91 (t, J = 7.4 Hz, 3H, CH₃), 1.31 (m, 2H, CH₂), 1.62 (m, 2H, CH₂), 3.97 (m, 1H, CH), 7.43 (AA'XX', 2H, ArH), 7.58 (AA'XX', 2H, ArH)

Example 4*3-Fluoro-N-methoxy-N-methylbenzamide*

To a suspension of 3-fluorobenzoic acid (140 mg, 1 mmol) and *N,O*-dimethylhydroxylamine hydrochloride (107 mg, 1.1 mmol) in dichloromethane (2.5 mL) was added 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (EDC) (211 mg, 1.1 mmol) and the mixture stirred at room temperature for 75 h. The solvents were removed under reduced pressure and the residue chromatographed using ethyl acetate-hexane (4:6) to separate the pure product (130 mg, 71%).

¹H-n.m.r. (CDCl₃) δ 3.36 (s, 3H, N-Me), 3.55 (s, 3H, N-OMe), 7.1-7.2 (m, 1H, ArH), 7.3-7.5 (m, 3H, Ar)

Example 5

141590721

44.

1-(3-Fluorophenyl)butan-1-one

To a solution of 3-fluoro-*N*-methoxy-*N*-methylbenzamide (130 mg, 0.71 mmol) in dry THF (2 mL) cooled to -10°C was added propyl magnesium chloride (532 μL , 2M solution in ether, 1.1 mmol) under nitrogen. The solution was stirred at -10°C for 1h and at room temperature for 75 min. The solution was the poured into saturated aqueous ammonium chloride and the product extracted into ethyl acetate (3 x 25 mL). The organic layers were combined, washed with brine, dried (MgSO_4) and concentrated to give a pale yellow oil, which was purified by chromatography using ethyl acetate-hexane (1:9) to separate the pure product (78 mg, 66%).

^1H -n.m.r. (CDCl_3) δ 1.01 (t, 3H, $J=7.4$ Hz, Me), 1.77 (m, 2H, CH_2CH_3), 2.93 (t, $J=7.2$ Hz, 2H, COCH_2), 7.15-7.30 (m, 1H, Ar), 7.35-7.50 (m, 1H, Ar), 7.60-7.70 (m, 1H, Ar), 7.70-7.80 (m, 1H, Ar).

Example 6

15 *1-(5-Methylpyridin-3-yl)butan-1-amine*

To a solution of 1-(5-methylpyridin-3-yl)butan-1-one (800 mg, 4.9 mmol) and ammonium formate (1.55 g, 24.5 mmol) in methanol (5 mL) under nitrogen was added dichloro(pentamethylcyclopentadienyl)rhodium (III) dimer (45 mg, 0.074 mmol). The solution was heated at reflux for 8 h. after which time the solution was cooled to room temperature and acidified to pH ~ 2 with 2M HCl. The mixture was washed with dichloromethane (3 x 15 mL) and the aqueous phase then basified to pH ~ 12 by addition of solid KOH. The aqueous phase was extracted with dichloromethane (3 x 15 mL) and

141590721

45.

the combined organic layers washed with brine, dried (Na_2SO_4) and concentrated to give a pure product (620 mg, 77%).

^1H -n.m.r. (CDCl_3) δ 0.91 (t, 3H, $J=7.4$ Hz, $\text{CH}_2\text{CH}_2\text{CH}_3$), 1.14-1.82 (m, 6H, $2 \times \text{CH}_2$, NH), 2.33 (s, 3H, CH_3), 3.91 (t, 1H, $J=6.8$ Hz, CHNH_2), 7.47 (s, 1H, ArH), 8.32 (br s, 2H, Ar).

5 Example 7

(1R)-1-(4-Fluorophenyl)butan-1-ol



- Diethylaniline-borane complex (2.4 mL, 13.4 mmol) was added dropwise over 5 min. to a solution of (1S,2R)-1-aminoindan-2-ol (201 mg, 1.4 mmol) in dry THF (20 mL) cooled to 0°C under nitrogen. After 40 min. a solution of the ketone in THF (40 mL) was added dropwise over 90 min. The cooling bath was then removed and the solution then stirred at RT for 5h. Acetone (6 mL) was added and the resulting solution stirred at RT for 60 min. before the solvents were removed under reduced pressure. The residue was dissolved in toluene (100 mL) and washed successively with 1M H_2SO_4 (4 x 50 mL), H_2O (2 x 50 mL) and brine (50 mL) and then dried (Na_2SO_4).

^1H -n.m.r. (CDCl_3) δ 0.93 (t, 3H, $J=7.2$ Hz, $\text{CH}_2\text{CH}_2\text{CH}_3$), 1.17-1.51 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_3$), 1.58-1.82 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_3$), 4.66 (t, $J=6.6$ Hz, CHNH_2), 6.98-7.07 (m, 2H, ArH), 7.26-7.34 (m, 2H, ArH).

Example 8

- 20 (1R)-1-Pyridin-3-ylbutan-1-ol



To a solution of S-(-)- α,α -diphenyl-2-pyrrolidinemethanol (500 mg, 2 mmol) and trimethylborate (0.27 mL, 2.4 mmol) in THF (20 mL) was added borane-dimethylsulfide

141590721

46.

complex (20mL, 2M in THF, 40 mmol). The mixture was cooled to 0°C and 1-pyridin-3-ylbutan-1-one was added dropwise over 4 h. The solution was stirred at RT overnight and then treated with 2M HCl (175 mL) and stirring continued for 4 h. The volatiles were removed under reduced pressure and the resulting cloudy solution basified to pH 11 with aqueous ammonia. The product was extracted into ethyl acetate (3 x 100 mL) and the combined organic layers washed with brine (100 mL) and dried (Na₂SO₄). Removal of the solvent under reduced pressure afforded ~3g of product.

¹H-n.m.r. (CDCl₃) 80.94 (t, 3H, *J*=7.4 Hz, CH₂CH₂CH₃), 1.20-1.54 (m, 2H, CH₂CH₂CH₃), 1.60-1.91 (m, 2H, CH₂CH₂CH₃), 2.64 (br s, 1H, OH), 4.73 (t, 1H, *J*=6.6 Hz, CHOH), 7.24-7.30 (m, 1H, Ar), 7.68-7.74 (m, 1H, Ar), 8.47-8.52 (m, 2H, Ar).

Example 9

3-Amino-3-phenylpropan-1-ol




To a suspension of 3-amino-3-phenylpropanoic acid (2.0 g, 12.1 mmol) in dry THF (45 mL) cooled to 0 °C under N₂ was added portionwise over 20 minutes solid LiAlH₄ (920 mg, 24.2 mmol). Stirring was continued at room temperature for 24 h. after which time solid Na₂SO₄·10H₂O was added with stirring until only a heavy white precipitate was present. The organic layer was diluted with ether and filtered through Celite® and the concentrated *in vacuo*. The residue was dissolved in ethyl acetate (50 mL) and extracted with 1N HCl (3 x 40 mL). The aqueous layers were combined and basified to pH~12 with 5M NaOH. The aqueous phase was extracted with ethyl acetate (3 x 50 mL) and the combined organic phases dried (Na₂SO₄) and concentrated *in vacuo* to separate the product which was used without further purification (0.9g, 49%).

¹H-n.m.r. (CDCl₃) δ 1.84-1.94 (m, 2H, CH₂CH₂OH), 2.68 (br s, 1H, NH₂), 3.79 (t, 2H, *J*=5.8 Hz, CH₂CH₂OH), 4.08-4.15 (m, 1H, CHNH₂), 4.77 (q, 1H, CH), 7.21-7.38 (m, 5H, ArH)

Example 10

141590721

47.

6-Chloro-N-[1-(4-methylphenyl)butyl]pyrazin-2-amine

To a solution of 1-[4-(trifluoromethyl)phenyl]butan-1-amine (0.40 g, 1.9 mmol) and 2,6-dichloropyrazine (0.55 g, 3.7 mmol) in 1,4-dioxane (6 mL) was added anhydrous potassium carbonate (0.39 g, 2.8 mmol). The resulting mixture was heated at reflux for 18 h. After cooling to room temperature, the mixture was diluted with ethyl acetate and H₂O. The organic phase was collected, dried and concentrated. The residue was purified by flash chromatography eluting with ethyl acetate-hexane (1:1) to give pure product (0.03 g, 5%).

¹H-n.m.r. (CDCl₃) 80.95 (t, 3H, *J* = 7.4 Hz, CH₃), 1.39 (m, 2H, CH₂), 1.81 (m, 2H, CH₂), 4.77 (q, 1H, CH), 5.09 (br d, 1H, NH), 7.42-7.62 (m, 5H, ArH), 7.80 (s, 1H, pyraz-H)

Example 11*N-[3-[1-[(6-Chloropyrazin-2-yl)amino]butyl]phenyl]acetamide*

To a stirred solution of *N*-[1-(3-aminophenyl)butyl]-6-chloropyrazin-2-amine (0.10 g, 0.36 mmol) and triethylamine (100 μ L, 0.72 mmol) in dichloromethane (3 mL) cooled to 0 °C. was added acetyl chloride (31 μ L, 0.43 mmol). The resulting mixture was stirred at room temperature for 18 h. then diluted with dichloromethane (10 mL) and washed with H₂O (10 mL) and brine (10 mL). The organic phase was collected, dried and concentrated and the residue purified by flash chromatography eluting with ethyl acetate-hexane (1:1) to give pure product (94 mg, 82%).


141590721

48.

¹H-n.m.r. (CDCl₃) δ 0.93 (t, 3H, *J*=7.4 Hz, Me), 1.28-1.43 (m, 2H, CH₂), 1.76-1.85 (m, 2H, CH₂), 2.16 (s, 3H, COCH₃), 4.48-4.59 (m, 1H, CHCH₂CH₂CH₃), 5.15 (br d, *J*=6.8 Hz, 1H, NH), 7.06 (d, 1H, *J*=7.4 Hz, Ar), 7.23-7.33 (m, 3H, Ar), 7.54 (s, 1H, CONH), 7.60 (s, 1H, pyraz-H), 7.77 (s, 1H, pyraz-H).

5 Example 12

N-[2-Methoxy-4-(6-[(1*S*)-1-pyridin-3-ylbutyl]amino)pyrazin-2-yl]phenyl]-*N*-ethylurea



Under a nitrogen atmosphere a mixture of 6-chloro-*N*-[(1*S*)-1-pyridin-3-ylbutyl]pyrazin-2-amine (53 g, 0.2 mmol), 4-[[[(ethylamino)carbonyl]amino]-3-methoxyphenyl]boronic acid pinacol diester (69 mg, 0.23 mmol), tetrakis(triphenylphosphine)palladium(0) (23 mg, 0.02 mmol) in toluene-*n*-propanol (2.6 mL, 3:1) was treated with 2M aqueous sodium carbonate solution (0.15 mL, 0.3 mmol). The resulting mixture was stirred vigorously whilst being heated at 100°C for 22 hours. Once cool ethyl acetate (10 mL) was added and the mixture washed with H₂O (6 x 10 mL) and brine (10 mL) and then dried (Na₂SO₄). Removal of solvent *in vacuo* then yielded crude product, which was purified by column chromatography using dichloromethane-methanol-aqueous ammonia (93:7:1) as eluent to furnish the product (60 mg).


¹H-n.m.r. (CDCl₃) δ 0.98 (t, 3H, *J*=7.4 Hz, CH₃), 1.14 (t, 3H, *J*=7.2 Hz, CH₃), 1.36-1.52 (m, 2H, CH₂), 1.78-1.98 (m, 2H, CH₂), 2.27 (s, 3H, CH₃), 3.23-3.37 (m, 2H, CH₂), 4.84-4.94 (m, 2H, CH, NHCONH), 5.19 (d, 1H, *J*=6.2 Hz, CHNH), 6.31 (s, 1H, ArNH), 7.24-7.30 (m, 1H, ArH), 7.55-7.73 (m, 4H, ArH), 7.75 (s, 1H, pyraz-H), 8.21 (s, 1H, pyraz-H), 8.49-8.52 (m, 1H, ArH), 8.69 (d, 1H, *J*=2.0 Hz, ArH).

Example 13

4-[[[(Ethylamino)carbonyl]amino]-3-methylphenyl]boronic acid pinacol diester

141590721

49.



The aniline (500mg, 2.7 mmol) was dissolved in pyridine (5 mL) and to this was added ethyl isocyanate (424 μ L, 5.4 mmol). The resulting solution was stirred overnight during which time a thick precipitate formed which was filtered and dried *in vacuo* to give the ethyl urea (595 mg, 86%). ¹H-n.m.r. (d_6 -DMSO) δ 1.05 (t, 3H, J =7.4 Hz, CH₃), 2.15 (s, 3H, CH₃), 3.09 (dq, 2H, J =7.4, 5.4 Hz, CH₂), 6.55 (br t, 1H, J =5.4 Hz, NH), 7.24 (dd, 1H, J =8.8, 2.2 Hz, Ar-H), 7.30 (d, 1H, J =2.2 Hz, Ar-H), 7.66 (s, 1H, NH), 7.82 (d, 1H, J =8.8 Hz, Ar-H).

To a degassed suspension of the ethyl urea bromide (514 mg, 2 mmol), potassium acetate (784 mg, 8 mmol) and bis(pinacolato)diboron (508 mg, 2 mmol) in a mixture of ethanol (10 mL) and dioxane (2 mL) was added dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium (II) dichloromethane adduct (44 mg, 0.06 mmol). The solution was heated at reflux for 2 h. and then allowed to cool to RT. The solvent was removed under reduced pressure and the residue dissolved in dichloromethane (60 mL). This was washed with H₂O (2 x 30 mL) and brine (30 mL), dried (Na₂SO₄) and concentrated *in vacuo*. The residue was chromatographed using ethyl acetate-hexane (1:1) as eluant to separate the pure product as a solid (351 mg, 58%).

¹H-n.m.r. (CDCl₃) δ 1.12 (t, 3H, J =6.8 Hz, CH₃), 1.34 (s, 12H, CH₃), 2.26 (s, 3H, CH₃), 3.09 (dq, 2H, J =6.8, 5.2 Hz, CH₂), 4.69 (br t, 1H, NH), 6.06 (s, 1H, NH), 7.49 (d, 1H, J =7.8 Hz, Ar-H), 7.67 (m, 2H, Ar-H).

Example 14

5-Bromo-N-(1-phenylbutyl)pyridin-3-amine



141590721


50.

To a degassed solution of 3,5-dibromopyridine (0.4 g, 1.7 mmol), sodium *tert*-butoxide (227 mg, 3.6 mmol), tris(dibenzylideneacetone)dipalladium (145 mg, 0.09 mmol) and *rac*-BINAP (105 mg, 0.2 mmol) in dry toluene (10 mL) was added 1-phenylbutan-1-amine (0.2 mL). The mixture was heated at reflux for 24h and upon cooling to room temperature
5 was diluted with ether and washed with brine (3 x 30 mL). The solution was dried (Na_2SO_4) and concentrated and the residue chromatographed using ethyl acetate-hexane (40:60) as eluant. From the early fractions 97 mg of pure product was obtained.

^1H -n.m.r. (CDCl_3) δ 0.94 (t, 3H, $J = 7.2$ Hz, CH_3), 1.21-1.53 (m, 2H, CH_2), 1.68-1.87 (m, 2H, CH_2), 4.19-4.32 (m, 2H, CH and NH), 6.86 (t, 1H, $J = 2.0$ Hz, ArH), 7.20-7.38 (m, 5H, ArH),
10 7.88 (d, 1H, $J = 2.0$ Hz, ArH), 7.92 (d, 1H, $J = 2.0$ Hz, ArH).

Example 15

5-[4-(Benzyloxy)-3-methoxyphenyl]-3-chloro-6-methyl-1,2,4-triazine



A degassed solution of 3,5-dichloro-6-methyl-1,2,4-triazine (151 mg, 0.9 mmol), 4-
15 (benzyloxy)-3-methoxybenzene boronic acid (236 g, 0.9 mmol), tetrakis(triphenylphosphine)palladium(0) (55 mg, 0.05 mmol) and tripotassium phosphate (249 mg, 1.2 mmol) in dimethoxyethane (5 mL) was heated at reflux for 24 h. Upon cooling to room temperature the solution was diluted with CHCl_3 and filtered through Celite®. The filtrate was concentrated *in vacuo* and the product purified by column
20 chromatography using ethyl acetate-hexane (25:75) as eluent to furnish (75 mg, 22%).


^1H -n.m.r. (CDCl_3) δ 2.88 (s, 3H, Ar- CH_3), 3.98 (s, 3H, OCH_3), 5.26 (s, 2H, CH_2), 6.99 (d, 1H, $J = 8.5$ Hz, ArH), 7.25-7.44 (m, 7H, ArH).

Example 16

5-[4-(Benzyloxy)-3-methoxyphenyl]-6-methyl-N-(1-phenylpropyl)-1,2,4-triazin-3-amine

141590721

51.




A solution of the triazine (50 mg, 0.14 mmol) in ethoxyethanol (2.5 mL) was treated with diisopropylamine (50 μ L, 0.28 mmol) and 1-phenylbutan-1-amine (46 μ L, 0.28 mmol) and the resulting solution heated at 110°C for 3 days. The solvent was then removed *in vacuo* to give an orange residue which was chromatographed using ethyl acetate-dichloromethane (20:80) as eluant to separate the product as a pale yellow solid (60 mg, 90%).

¹H-n.m.r. (CDCl₃) δ 0.94 (t, 3H, CH₃), 1.21-1.53 (m, 2H, CH₂), 1.78-1.97 (m, 2H, CH₂), 2.62 (s, 3H, Ar-CH₃), 3.89 (s, 3H, OCH₃), 5.10 (br q, 1H, CH₂), 5.22 (s, 2H, CH₂), 5.65 (br s, 1H, NH), 6.87 (d, 1H, *J* = 9.2 Hz, ArH), 7.12-7.44 (m, 12H, Ar-H).

Example 17

2-Chloro-5-methyl-N-[(1*S*)-1-pyridin-3-ylbutyl]pyrimidin-4-amine



To a solution of (1*S*)-1-pyridin-3-ylbutan-1-amine (600 mg, 4.0 mmol) in ethanol (20 mL) was added 2,4-dichloro-5-methylpyrimidine (717 mg, 4.4 mmol) and diisopropylethylamine (1.4 mL, 8mmol). The solution was heated at reflux for 24h after which time the solvent was removed under reduced pressure. The residue was dissolved in ethyl acetate (30 mL) and washed with H₂O (2 x 15 mL), brine (15 mL) and dried (Na₂SO₄). The residue remaining after concentration *in vacuo* was chromatographed using ethyl acetate-hexanes (9:1) as eluant to separate the desired product (370 mg).

¹H-n.m.r. (CDCl₃) δ 0.96 (t, 3H, *J* = 7.2 Hz, CH₃), 1.23-1.52 (m, 2H, CH₂), 1.85-1.99 (m, 2H, CH₂), 2.03 (d, 3H, *J* = 0.8 Hz, Ar-Me), 5.03 (br d, 1H, *J* = 7.2 Hz, NH), 5.22-5.33 (m, 1H,


141590721

52.

CH), 7.24-7.30 (m, 1H, ArH), 7.65-7.71 (m, 1H, ArH), 7.81 (d, 1H, $J=0.8$ Hz, ArH), 8.51 (dd, $J=5.6, 1.4$ Hz, 1H, ArH), 8.64 (d, 1H, $J=1.8$ Hz, ArH).

Example 18

N-Ethyl-*N*'-[2-methoxy-4-(5-methyl-4-[(*1S*)-1-pyridin-3-ylbutyl]amino)pyrimidin-2-yl)phenyl]urea




Under a nitrogen atmosphere a mixture of 2-chloro-5-methyl-*N*-[(*1S*)-1-pyridin-3-ylbutyl]pyrimidin-4-amine (277 mg, 1.0 mmol), 4-[[[(ethylamino)carbonyl]amino]-3-methoxyphenyl]boronic acid pinacol diester (416 mg, 1.3 mmol), tetrakis(triphenylphosphine)palladium(0) (116 mg, 0.1 mmol) in toluene-*n*-propanol (12 mL, 3:1) was treated with 2M aqueous sodium carbonate solution (750 μ L, 1.5 mmol). The resulting mixture was stirred vigorously whilst being heated at 100 °C for 17 hours. Once cool ethyl acetate (25 mL) was added and the mixture washed with H₂O (6 x 15 mL), brine (20 mL) and dried (Na₂SO₄). Removal of solvent *in vacuo* then yielded crude product, which was purified by column chromatography using dichloromethane-methanol-aqueous ammonia (93:7:1) as eluent to furnish the product (110 mg, 57%).

¹H-n.m.r. (CDCl₃) δ 0.99 (t, 3H, $J=7.4$ Hz, CH₃), 1.18 (t, 3H, $J=7.2$ Hz, CH₃), 1.33-1.60 (m, 2H, CH₂), 1.82-2.02 (m, 2H, CH₂), 2.11 (s, 3H, Ar-Me), 3.25-3.39 (m, 2H, CH₂), 3.87 (s, 3H, OMe), 4.80-4.96 (m, 2H, 2 x NH), 5.26-5.36 (m, 1H, CH), 6.98 (br s, 1H, Ar-NHCONH), 7.22-7.28 (m, 1H, ArH), 7.67-7.72 (m, 2H, ArH), 7.83-7.88 (m, 1H, ArH), 8.05 (d, 1H, $J=0.8$ Hz, ArH), 8.10 (d, 1H, $J=8.2$ Hz, ArH), 8.47 (dd, 1H, $J=6.6, 1.8$ Hz, ArH), 8.69 (d, 1H, $J=2.0$ Hz, ArH).

Example 19

141590721


53.

6-Chloro-N-(1-pyridin-3-ylbutyl)pyrimidin-4-amine

To a solution of 1-pyridin-3-ylbutan-1-amine (100 mg, 0.67 mmol) in ethanol (4 mL) was added 4,6-dichloropyrimidine (109 mg, 0.73 mmol) and diisopropylethylamine (232 μ L, 1.33 mmol). The solution was stirred for 3 days after which time the solvent was removed under reduced pressure. The residue was dissolved in ethyl acetate (25 mL) and washed with H_2O (3 x 15 mL), brine (15 mL) and dried (Na_2SO_4). The residue remaining after concentration in vacuo was chromatographed using ethyl acetate-hexanes (4:6 - 9:1) as eluant to separate the desired product (28 mg).

1H -n.m.r. ($CDCl_3$) δ 0.96 (t, 3H, $J=7.2$ Hz, CH_3), 1.22-1.51 (m, 2H, CH_2), 1.76-1.96 (m, 2H, CH_2), 4.76 (br s, 1H, CH), 5.63 (br s, 1H, NH), 6.23 (s, 1H, ArH), 7.26-7.32 (m, 1H, ArH), 7.59-7.65 (m, 1H, ArH), 8.33 (s, 1H, ArH), 8.53-8.56 (m, 1H, ArH), 8.60 (d, 1H, $J=1.8$ Hz, ArH).

Example 20

15 *2-Methoxy-4-[6-[(1-pyridin-3-ylbutyl)amino]pyrimidin-4-yl]phenol*

Using an analogous procedure to that described in Example 12, reaction of 6-chloro-N-(1-pyridin-3-ylbutyl)pyrimidin-4-amine (24 mg, 0.09 mmol) with 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenol (27 mg, 0.11 mmol) afforded the desired product as a brown foam (12 mg).

141590721

54.

¹H-n.m.r. (CDCl₃) δ 0.97 (t, 3H, *J*=7.2 Hz, CH₃), 1.32-1.55 (m, 2H, CH₂), 1.83-1.95 (m, 2H, CH₂), 3.94 (s, 3H, OMe), 4.87 (br s, 1H, CH), 5.43 (br s, 1H, NH), 6.50 (s, 1H, ArH), 6.93 (d, 1H, *J*=8.4 Hz, ArH), 7.25-7.31 (m, 2H, ArH), 7.55 (d, 1H, *J*=2.0 Hz, ArH), 7.66-7.71 (m, 1H, ArH), 8.53 (dd, 1H, *J*=4.9, 1.6 Hz, ArH), 8.58 (d, 1H, *J*=1.6 Hz, ArH), 8.66 (d, 1H, *J*=2.0 Hz, ArH).

5

141590721

55.

SCREENING

Compound Dilution

For screening purposes, compounds were diluted in 96 well plates at a concentration of 20 μ M. Plates were warmed at 37°C for 30 minutes before assay.

5 Growth and Maintenance of Cancer Cell lines

K562 (Chronic Myeloid Leukemia), PC3 (Prostate Cancer), and DU145 (Prostate Cancer) were obtained from the American Type Culture Collection (ATCC). K562 was grown in RPMI, with 10% FBS with Glutamax added. DU145 cells were cultured in DMED, with 10% FBS and Glutamax and MEM non-essential amino acids added. PC3 cells were grown in F12K medium, with 10% FBS and Glutamax and MEM non-essential amino acids added. All cells were grown at 37°C in 5% CO₂.

Establishment of TEL:JAK cell lines

The coding region encompassing nucleotides 1-487 of TEL was amplified by PCR using the oligonucleotides 5TEL (5'-GGA GGA TCC TGA TCT CTC TCG CTG TGA GAC-3') and 3TEL (5'-AGGC GTC GAC T'IC T'IC T'IC ATG GTT CTG-3') and U937 mRNA as template. A BamHI site was present into the 5TEL Primer, a Sal I site was incorporated into the 3TEL primer. The regions encompassing the kinase domains of JAK2 (nucleotides 2994-3914; ; JAK2F 5'-ACGC GTC GAC GGT GCC TTT GAA GAC CCG GAT-3' ; JAK2R 5'-ATA GTT TAG CCG CCG C'IC AGA ATG AAG GTC ATT T-3') and JAK3 (nucleotides 2520-3469; JAK3F 5'-GAA GTC GAC TAT GCC TGC CAA GAC CCC ACG ATC TT-3' ; JAK3R 5'-GGA TCT AGA CTA TGA AAA GGA CAG GGA GTG GTG TTT -3') were generated by PCR using Taq DNA Polymerase (Gibco/BRL) and U937 mRNA as template. A SalI site was incorporated into the forward primer of JAK2 and JAK3, a Not I site was incorporated into the ~~JAK2~~ reverse primer and a Xba I site was added to the reverse primer of JAK3.

A TEL/Jak2 fusion was generated by digestion of the TELPCR product with BamHI /Sal I, digestion of the JAK2 PCR product with Sal I/ Not I followed by ligation and subcloning into the mammalian expression Vector pTRE 2 (Clontech) digested with BamHI-Not I (pTELJAK2). For JAK3 Sal I/ Not I cleaved kinase domain PCR product

141590721

56.

was ligated with BamHI / Sal I cleaved TELproduct followed by ligation into BamHI / Not I cleaved pTRE2 (pTELJAK3).

- 5 The growth factor dependent myelomonocytic cell line BaF3 bearing the pTET-off plasmid (Clontech) was transfected with either pTELJAK2 or pTELJAK3 and the cells selected for factor independent growth. BaF 3 wild type cells were cultured in DMEM 10% FCS, 10% WEHI 3B conditioned medium. BaF3 TELJAK cells were cultured in DMEM 10% Tet-System Approved FBS (without WEHI 3B conditioned medium).

Cellular assays were performed as follows:

- 10 Cell suspensions were prepared by harvesting cells from culture. (Cells used in this test should be in later log phase growth and high viability.) Cells were diluted in correct growth medium to 1.1x final concentration (from 50000 cell/mL to 200,000 cell/mL, depending on cell line).

- 15 Compounds to be tested were added (10µL, 10X final concentration) to a flat bottom 96-well plate. The cellular suspension (90µL per well) was added, and the plate incubated for 40 hr at 37 °C, 5% CO₂. MTT (20 µL per well, 5mg/mL in PBS) was added and the plates were returned to the incubator for a further 6 hours. Lysis buffer (100 µL per well, 10% SDS, 0.01N HCl) was added and the plate stored in the incubator overnight. The plate was then read at 590 nm.

Tubulin Assay

- 20 Turbidometric assays of microtubule assembly were performed by incubating microtubule protein in cuvettes at 37 °C in a thermostatically controlled spectrophotometer measuring the change in absorbance at 340nm over time. The microtubule protein was incubated with the each test compound at 0 °C and polymerisation was initiated by addition of 1mM GTP, prior to heating to 37 °C.

25 Results

The activity of a range of compounds is shown in Tables 1 and 2. Compounds that exhibited a capacity to inhibit greater than 50% of cellular growth at a concentration of 20µM are designated as "+"; compounds which did not inhibit 50% of cellular growth at

141590721







57.

20 μ M are designated as "-"; compounds which were not tested are designated as "NT".
Likewise, compounds which inhibited tubulin polymerisation by greater than 50% at
50 μ M are designated as "+"; compounds which did not inhibit tubulin polymerisation by
50% at 50 μ M are designated as "-"; and compounds which were not tested are designated
5 as "NT".

141590721







58.

Table 1

CHEMISTRY	DU145	PC3	K562	TelJak2	TelJak3	Tubulin
 C24H28FN5O2	+	+	+	+	+	+
 C27H37N5O2	-	-	+	+	+	NT
 C21H23N3O	+	+	+	+	+	+
 C26H32N4	-	-	-	+	+	NT
 C24H29N3O3	+	+	+	+	+	+
 C27H36N4O2	+	+	+	+	+	+







141590721

59.

CHEMISTRY	DU145	PC3	K562	TelJak2	TelJak3	Tubulin
 C24H29N5O2	+	+	+	+	+	+
 C25H30N4O	-	-	-	+	-	NT
 C26H33N5	-	-	-	+	+	NT
 C25H28N6O2	+	+	+	+	+	+
 C25H31N3O3	+	+	+	+	+	+
 C26H35N4O3	-	+	+	+	+	NT







141590721

60.

CHEMISTRY	DU145	PC3	K562	TelJak2	TelJak3	Tubulin
 C21H24N4	+	+	+	+	+	+
 C26H33N7O2	-	+	-	+	+	NT
 C28H35N4O2	+	+	+	+	+	+
 C28H38N4O2	+	+	+	+	+	+
 C28H39N5O2	+	+	+	+	+	+
 C27H31N5O2	+	+	+	+	+	+







141590721

61.

CHEMISTRY	DU145	PC3	K562	TelJak2	TelJak3	Tubulin
 C26H28N5O2	-	-	+	+	+	NT
 C27H33N6O	+	+	+	+	+	+
 C28H38N4O3	-	-	+	+	+	NT
 C30H40N4O2	+	+	+	+	+	+
 C29H38N4O2	+	+	+	+	+	+
 C28H40N4O2	+	+	+	+	+	+







141590721

62.

CHEMISTRY	DU145	PC3	K562	TelJak2	TelJak3	Tubulin
 C30H41NSO2	+	+	+	+	+	+
 C25H32N4O2	+	+	+	+	+	+
 Chemistry 548	-	-	+	+	+	NT
 C28H32N4O	-	-	+	+	+	NT
 C26H32N4O	-	-	+	+	+	NT
 C22H28N8O	-	-	+	+	+	NT







141590721

63.

CHEMISTRY	DU145	PC3	K562	TelJak2	TelJak3	Tubulin
 C26H26N5O2	-	-	+	+	+	NT
 C27H33N5O	+	+	+	+	+	+
 C21H24N4O2	+	+	+	+	+	+
 C21H24N4O3	-	-	-	+	+	NT
 C25H30N4	+	+	+	+	+	+
 C21H26N5O	+	+	+	+	+	+







141590721

64.

CHEMISTRY	DU145	PC3	K562	TelJak2	TelJak3	Tubulin
 G22H26N6O2	+	+	+	+	+	+
 G19H18N3O2	+	+	+	+	+	+
 G24H23N2O3	-	+	+	+	+	NT
 G20H20N3O2	+	+	+	+	+	+
 G22H24N2O2	+	+	+	+	+	+
 G23H28N2O	+	+	+	+	+	+







141590721

65.

CHEMISTRY	DU145	PC3	K562	TelJak2	TelJak3	Tubulin
 C26H32N8O2	-	-	+	+	-	NT
 C25H31N5O	-	-	-	+	+	NT
 C28H37N5O	+	-	+	+	+	NT
 C28H35N5O2	-	-	-	+	+	NT
 C26H31N5O	NT	-	-	+	+	NT
 C25H28N4O2	NT	-	-	+	+	NT







141590721

66.

CHEMISTRY	DU145	PC3	K562	TelJak2	TelJak3	Tubulin
 C21H24N6O2	NT	-	NT	+	+	NT
 C23H28N6O	+	+	+	+	+	+
 C28H38N6O2	-	-	-	+	+	NT
 C27H35N5O2	-	-	+	+	+	NT
 C27H35N5O3	-	-	-	+	+	NT
 C28H37N7O3	+	+	+	+	+	+







141590721

67.

CHEMISTRY	DU145	PC3	K562	TelJak2	TelJak3	Tubulin
 C25H39N7O2	+	+	+	+	+	+
 C28H39N7O2	-	-	-	+	+	NT
 C24H30N6O	+	+	+	+	+	+
 C24H30N6O2	+	+	+	+	+	+
 C23H28N6O2	+	+	+	+	+	+
 C23H28N6O	+	+	+	+	+	+






141590721

68.

CHEMISTRY	DU145	PC3	K562	TelJak2	TelJak3	Tubulin
 C21H24N4O2	-	-	+	+	+	NT
 C23H28N6O	+	+	+	+	+	+
 C23H29N6O2	+	+	+	+	+	+
 C24H30N6O2	+	+	+	+	+	+
 C24H30N6O	+	+	+	+	+	+
 C24H30N6O2	+	+	+	+	+	NT







141590721

69.

CHEMISTRY	DU145	PC3	K562	TelJak2	TelJak3	Tubulin
 C23H28N8O2	+	+	+	+	+	+
 C24H30N8O2	+	+	+	+	+	+
 C26H34N6O	-	-	+	+	+	NT
 C22H26N6O2	+	+	+	+	+	+
 C25H31N5O2	+	+	+	+	+	+







141590721

70.

CHEMISTRY	DU145	PC3	K562	TelJak2	TelJak3	Tubulin
 C24H28N5O3	+	+	+	+	+	+
 C23H28N6O2	+	+	+	+	+	+
 C21H23N3O3	+	-	+	+	+	NT
 C23H27N5O3	+	+	+	+	+	+
 C24H29N5O3	+	+	+	+	+	+
 C20H22N4O2	+	+	+	+	+	+







141590721

71.

CHEMISTRY	DU145	PC3	K562	TelJak2	TelJak3	Tubulin
 C24H30N8O2	+	+	+	+	+	+
 C21H24N4O2	+	+	+	+	+	+
 C23H28N8O2	+	+	+	+	+	+
 C20H22N4O2	+	+	+	+	+	+
 C23H28N4O3	-	-	+	+	+	NT
 C18H20N4O2	-	-	+	+	+	NT



141590721

72

CHEMISTRY	DU145	PC3	K562	TelJak2	TelJak3	Tubulin
 C24H30N6O2	+	+	+	+	+	+
 C21H24N4O2	+	+	+	+	+	+
 C23H28N6O2	+	+	+	+	+	+
 C20H22N4O2	+	+	+	+	+	+
 C23H28N4O3	-	-	+	+	+	NT
 C19H20N4O2	-	-	+	+	+	NT

141590721







73.

CHEMISTRY	DU145	PC3	K562	TelJak2	TelJak3	Tubulin
 C23H28N8O2	+	+	+	+	+	+
 Q19H19N3O2	+	+	+	+	+	+

141590721

74.

Table 2

Chemistry	TelJak2	TelJak3	DU145	Tubulin
 C25H23N5O3	+	+	+	+
 C24H22N4O2	+	+	+	+
 C24H27N5O3	+	+	+	+
 C25H23N5O3	+	+	+	+
 C20H21N3O2	+	+	+	+
 C19H19N3O2	+	+	+	+

141590721

. 75

It will be appreciated by persons skilled in the art that numerous variations and/or modifications may be made to the invention as shown in the specific embodiments without departing from the spirit or scope of the invention as broadly described. The present embodiments are, therefore, to be considered in all respects as illustrative and not restrictive.

Dated this 3rd day of December, 2003

Cytopia Pty Ltd

Patent Attorneys for the Applicant:

**BLAKE DAWSON WALDRON PATENT
SERVICES**

141590721

76.

References

Jordan M A. and Wilson L. (1998) Microtubules and actin filaments: dynamic targets for cancer chemotherapy. *Curr. Op. Cell Biol.*, 10, 123-130.

Jordan MA, Margolis RL, Himes RH, Wilson L., (1986) Identification of a distinct class of vinblastine binding sites on microtubules. *J. Mol. Biol.* 187:61-73

Rai SS, and Wolff J., (1996) Localization of the vinblastine-binding site on -tubulin. *J. Biol. Chem.* 271:14707-11

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/AU04/001689

International filing date: 03 December 2004 (03.12.2004)

Document type: Certified copy of priority document

Document details: Country/Office: AU
Number: 2003906680
Filing date: 03 December 2003 (03.12.2003)

Date of receipt at the International Bureau: 04 January 2005 (04.01.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record.**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☒ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.